# ## FIRST THINGS FIRST

La Paz/Mohave Region



2022 Needs and Assets Supplemental Report:

Children's Access to and Use of Public Health Services

## **Report Prepared by:**

Center for Health Information & Research
Arizona State University
502 E. Monroe St, Suite C320

Phoenix, AZ 85004

(602) 496-2009 | chir@asu.edu / chs.asu.edu/chir

## **Project Team**

- Molly Loughran, MS, Data Science Specialist
- · Varnika Angampally, MS, Statistical Programmer
- Nishanth Prathap, MS, Data Science Specialist
- Tameka Sama, MBA, CRA, Center Administrator
- Sruthi Kommareddy, Database Analyst
- Gevork Harootunian, MS, Data Science Consultant
- Meghan Morris, MD, Clinical Consultant
- Anita Murcko, MD, Clinical Consultant

## Acknowledgements

The authors gratefully acknowledge the cooperation of First Things First representatives Melissa Affronti, Evaluation Project Director, and Roopa Iyer, Senior Director for Research and Evaluation for their valuable technical support and for the feedback provided by the Regional Needs and Assets (RNA) Health Base Expansion Workgroup along with the RNA Workgroup. Special thanks to Candida Hunter and Kalvina Belin for their skilled administration and management of tribal relations and to Lisa Colling for her assistance with Geocoding and mapping of FTF regions and subregions.



# **TABLE OF CONTENTS**

List of Tables	4
List of Figures	6
Executive Summary	8
Introduction	16
The Importance of Early Childhood Health	16
Definitions	16
Approach	19
Reporting	20
Data Limitations	21
La Paz/Mohave Regional Results	22
Population and Demographics of Children Enrolled in AHCCCS	23
Health Care Workforce	29
Primary Care	33
Well-Child Visits	36
Screening for Lead Poisoning	41
Weight Assessment and Counseling	44
Developmental Screening and Delay	45
Behavioral Health	54
Vision	57
Hearing	63
Oral Health	70
Immunizations	77
Maternal Prenatal and Postpartum Care	80
Health Plan Performance	85
Conclusion	89
References	90
Annendiy: Data Sources	95

## LIST OF TABLES

Table 1. Number of AHCCCS Children Birth to Age 5 by Year and Sex	. 23
Table 2. Number of AHCCCS Children Birth to Age 5 by Year and Age Group	. 23
Table 3. Number of AHCCCS Claims by Provider Type (Billing Entity), 2017-2019	. 28
Table 4. Supply of Key Health Professionals in Arizona per 1,000 AHCCCS Children, 2017-2019	.30
Table 5. Supply of Key Health Professionals in the La Paz/Mohave Region per 1,000 AHCCCS Childr	en,
2017-2019	.31
Table 6. Percent of AHCCCS Children Grouped by Travel Distance Between Provider and Child's Resident	nce
by Provider Type for Region and Arizona, 2017-2019	.32
Table 7. Arizona and Regional AHCCCS Rates for PCP Visits, 2017-2019	.33
Table 8. Percent of AHCCCS Children Ages 1 – 5 with a Visit to PCP by Subregion, 2017-2019	. 34
Table 9. Arizona and Regional AHCCCS Rates for Well-Child Visits, 2017-2019	.36
Table 10. Percent of AHCCCS Children Who Had Six or More Well-Child Visits During Their First 15 Mon	ıths
of Life by Subregion, 2017-2019	. 37
Table 11. Percent of AHCCCS Children Ages 3-5 with a Well-Child Visit by Subregion, 2017-2019	. 39
Table 12. Arizona and Regional AHCCCS Rates for Lead Poisoning Screening, 2017-2019	.41
Table 13. Percent of AHCCCS Children Who Had One or More Tests for Lead Poisoning by Their Second	ond
Birthday by Subregion, 2017-2019	. 42
Table 14. Arizona AHCCCS Rates for Weight Assessment and Counseling, Ages Birth to 5, 2017-2019	. 44
Table 15. National Medicaid HEDIS Rates for Weight Assessment and Counseling, Ages 3-17 Years, 20	17-
2019	. 45
Table 16. Arizona and Regional AHCCCS Rates for Developmental Screenings and Delay, 2017-2019	. 47
Table 17. Percent of Claims by Provider Type for AHCCCS Children Receiving Developmental Screening	ngs,
2017-2019	
Table 18. Percent of AHCCCS Children Receiving Developmental Screenings, Age 1-2, by Subregion, 20	
2019	
Table 19. Percent of AHCCCS Children with a Diagnosed Developmental Delay by Subregion, 2017-20	
Table 20. Percent of AHCCCS Children with a Diagnosed Developmental Delay Who Have Received	
Behavioral Health Services by Subregion, 2017-2019	
Table 21. Percent of Claims by Provider Type for AHCCCS Children with a Diagnosed Developmental De	
Who Have Received Behavioral Health Services, 2017-2019	
Table 22. Arizona and Regional AHCCCS Rates for Behavioral Health Services, Ages 3-5, 2017-2019	
Table 23. Percent of AHCCCS Children Ages 3-5 Receiving Behavioral Health Services by Subregion, 20	
2019	
Table 24. Percent of Claims by Provider Type for AHCCCS Children Ages 3-5 Receiving Behavioral Hea	
Services, 2017-2019	
Table 25. Arizona and Regional AHCCCS Rates for Vision, 2017-2019	
Table 26. Percent of AHCCCS Children Receiving Vision Screening or Well-Child Visit by Subregion, 20	
2019	
Table 27. Percent of AHCCCS Children Receiving Eye Exams by Subregion, 2017-2019	. 61

Table 28. Percent of AHCCCS Statewide and Regional Hearing Results, 2017-2019	64
Table 29. Percent of AHCCCS Children Under Age One Who Had Audiology Services of All Children	ı by
Subregion, 2017-2019	64
Table 30. Percent of Claims by Provider Type for AHCCCS Children Under Age One Who Had Audiol Services, 2017-2019	
Table 31. Percent of Claims by Provider Type for AHCCCS Children Ages 1-5 Receiving Hearing Screer Tests, 2017-2019	_
Table 32. Percent of AHCCCS Children Ages 1-5 Receiving Hearing Screening Tests by Subregion, 20 2019	
Table 33. Percent of AHCCCS Children Ages 1-5 Screened for Hearing and Who Had Additional Audiol Services by Subregion, 2017-2019	
Table 34. Percent of Audiology Service Claims by Provider Type for AHCCCS Children Ages 1-5 Scree for Hearing and Who Had Additional Audiology Services, 2017-2019	
Table 35. Percent of Statewide and Regional AHCCCS Children Oral Health Visits for Ages 1-5, 2017-2	
Table 36. AHCCCS Contractor Rate of Performance on Annual Dental Visits for Ages Two to 20 Years, 20	
Table 37. Percent of AHCCCS Claims by Provider Type for Children Ages 1-5 With at Least One Anr Dental Visit, 2017-2019	nual
Table 38. Percent of AHCCCS Children Ages 1-5 With at Least One Annual Dental Visit by Subregion, 20	017-
Table 39. Percent of Claims by Provider Type for AHCCCS Children Ages 1-5 Receiving Fluoride Varn 2017-2019	nish,
Table 40. Percent of AHCCCS Children Ages 1-5 Receiving Fluoride Varnish by Subregion, 2017-2019	75
Table 41. AHCCCS Statewide Aggregate Immunization Completion Rates by Two Years Old, FFY 2016	78
Table 42. Percent of Statewide and Regional AHCCCS Children Immunization Status, from AHCCCS Cla Data Only, 2017-2019	
Table 43. Percent of All AHCCCS Women Who Received Timely Prenatal and Postpartum Care, 2017-2	019
Table 44. Percent of Deliveries That Received a Prenatal Care Visit While Enrolled in AHCCCS in the F Trimester, on the Enrollment Start Date or Within 42 Days of Enrollment in AHCCCS by Subregion, 20	First 017-
2019 Table 45. Percent of AHCCCS Deliveries That Had a Postpartum Visit After Delivery by Subregion, 20 2019	017-
Table 46. AHCCCS Statewide Aggregate Health Plan Performance and Minimum Performance Standa 2017-2019	rds,
Table 47. Select Regional Indicators by AHCCCS Health Plan, 2017-2019	
Table 48. Select AHCCCS Statewide Indicators by Health Plan, 2017-2019	
Table 49. AHCCCS Complete Care Health Plans by Geographic Service Area	95

## LIST OF FIGURES

Figure 1. Number of AHCCCS Children Birth to Age 5 by Year and Subregion24	4
Figure 2. Number of AHCCCS Children Birth to Age 5 by Year and Race2	5
Figure 3. Number of AHCCCS Children Birth to Age 5 by Year and Ethnicity2	5
Figure 4. Number of AHCCCS Children Birth to Age 5 by Year and Tribal Affiliation20	6
Figure 5. Number of AHCCCS Children by Year and Health Plan2	7
Figure 6. Percent of AHCCCS Children Ages $1-5$ with a Visit to PCP by Age Group, Tribal Affiliation	
Ethnicity, Race and Year3	5
Figure 7. Percent of Regional AHCCCS Children by Number of Well-Child Visits Completed During Thei	ir
First 15 Months by Number of Visits and Year3	7
Figure 8. Percent of AHCCCS Children Who Had Six or More Well-Child Visits During Their First 15 Month	S
of Life by Race, Ethnicity, Tribal Affiliation and Year	8
Figure 9. Percent of AHCCCS Children Ages 3-5 with a Well-Child Visit by Race, Ethnicity, Tribal Affiliation	n
and Year40	0
Figure 10. Percent of AHCCCS Children Who Had One or More Tests for Lead Poisoning by Their Second	d
Birthday by Race, Ethnicity, Tribal Affiliation and Year4	3
Figure 11. Percent of AHCCCS Children Ages 3-5 Who Had Weight Assessment and Counseling by Year 4	5
Figure 12. Percent of AHCCCS Children Receiving Developmental Screenings by Age Group, Triba	
Affiliation, Ethnicity, Race and Year49	9
Figure 13. Percent of AHCCCS Children with a Diagnosed Developmental Delay by Ethnicity, Sex, Triba	ìε
Affiliation, Age Group and Year5	1
Figure 14. Percent of AHCCCS Children with a Diagnosed Developmental Delay Who Have Received	d
Behavioral Health Services by Ethnicity, Race, Sex, Tribal Affiliation, Age Group and Year5	3
Figure 15. Percent of AHCCCS Children Ages 3-5 Receiving Behavioral Health Services by Sex, Triba	
Affiliation, Ethnicity, Race and Year5	6
Figure 16. Percent of AHCCCS Children Receiving Vision Screening or Well-Child Visit by Age Group, Triba	϶l
Affiliation, Ethnicity, Race and Year60	0
Figure 17. Percent of AHCCCS Children Receiving Eye Exams by Age Group, Tribal Affiliation, Ethnicity	
Race and Year6	2
Figure 18. Percent of AHCCCS Children Under 1 Year of Age Who Had Audiology Services by Sex, Triba	϶l
Affiliation, Ethnicity and Year6	5
Figure 19. Percent of AHCCCS Children Ages 1-5 Receiving Hearing Screening Tests by Age Group, Triba	϶l
Affiliation, Ethnicity, Race, Sex and Year6	
Figure 20. Percent of AHCCCS Children Ages 1-5 Screened for Hearing and Who Had Additional Audiolog	
Services by Age Group, Ethnicity, Sex and Year6	
Figure 21. Percent of AHCCCS Children Ages 1-5 With at Least One Annual Dental Visit by Age Group, Triba	϶l
Affiliation, Ethnicity, Race and Year73	3
Figure 22. Percent of AHCCCS Children Ages 1-5 With One and Two Preventative Care Dental Visits in	
Year74	
Figure 23. Percent of AHCCCS Children Ages 1-5 Receiving Fluoride Varnish by Age Group, Tribal Affiliation	١,
Fthnicity, Race and Year 70	6

Figure 24. Percent of Deliveries That Received a Prenatal Care Visit While Enrolled in AH	CCCS in the First
Trimester, on the Enrollment Start Date or Within 42 Days of Enrollment in AHCCCS by A	ge Group, Triba
Affiliation, Ethnicity, Race and Year	82
Figure 25. Percent of AHCCCS Deliveries That Had a Postpartum Visit After Delivery by A	ge Group, Race
Ethnicity, Tribal Affiliation and Year	84

## **EXECUTIVE SUMMARY**

Under the direction of First Things First (FTF), the Arizona State University Center for Health Information & Research (CHiR) conducted a regional network analysis of children from birth to age 5 to determine the health assets and health needs in the La Paz/Mohave Region. The La Paz/Mohave Region consists of 11 subregions: Bullhead City, Colorado City, Dolan Springs, Fort Mohave, Kingman, Lake Havasu City, La Paz/Mohave Remainder, Littlefield, Parker, Quartzsite and Salome. The main data source was claims data from the Arizona Health Care Cost Containment System (AHCCCS), Arizona's Medicaid agency; therefore, the results presented in this report were for children and mothers who were enrolled in AHCCCS from 2017 to 2019. This population was denoted as AHCCCS children or AHCCCS women.

CHiR and representatives from the FTF Regions, Programs, and Evaluation teams determined priority indicators for this report. AHCCCS children's health was measured in the following categories: primary care and well-child visits, health care workforce, screening for lead poisoning, weight assessment and counseling, developmental health, behavioral health, vision, hearing, oral health, immunizations, maternal prenatal and postpartum care, and health plan performance. Many of the reported indicators were from the Healthcare Effectiveness Data and Information Set (HEDIS)<sup>2</sup>. HEDIS is a performance improvement tool whereby health plans, health care organizations and government agencies submit data on specific health measures. HEDIS uses the collected data to calculate national performance statistics and benchmarks and set standards for measures. HEDIS specifications were applied to the AHCCCS population for each region. Non-HEDIS indicators, which do not have associated benchmarks, were compared to state and national data when possible. The results were displayed by gender, age, race, ethnicity, tribal affiliation<sup>3</sup>, provider type, and health plan when the data was available and within data suppression guidelines. The results of the analyses are summarized below. When possible, the results are grouped by 1) indicators that met or were above the state average or national HEDIS standards and 2) indicators that did not meet or were below the state average or national HEDIS standards. Other notable findings are also presented that do not have comparison data.

## **Population and Demographics of Children Enrolled in AHCCCS**

There were 234,616 children from birth to age 5 enrolled in AHCCCS statewide from 2017 to 2019. In the La Paz/Mohave Region, there were 6,970 children enrolled in AHCCCS in 2017, 6,701 children enrolled in 2018 and 6,606 children enrolled in 2019. Of these, male AHCCCS children outnumbered females by 3-4%. There were 7% more infants and toddlers than preschoolers in 2017, with the difference dropping to

<sup>&</sup>lt;sup>1</sup> Data used in this report covers all AHCCCS members in Arizona, including members living in FTF tribal regions and subregions. Reports for tribal regions and subregions were carried out with specific approval from each tribe. For those tribal regions and subregions who did not give approval, data is included only in aggregate totals for Arizona, and—in the case of a tribal subregion—aggregate totals for the region.

<sup>&</sup>lt;sup>2</sup> See https://www.ncqa.org/hedis/

<sup>&</sup>lt;sup>3</sup> Tribal affiliation refers to whether an individual is a member of a federally recognized Arizona tribe and is displayed as a flag (Yes/No) in this report. This information is captured during enrollment in AHCCCS.

5% more in 2018 and 3% more in 2019. Approximately 87% of the AHCCCS children in the region lived in four subregions: Kingman (32-33%), Bullhead City (23%), Lake Havasu City (19-20%) and Fort Mohave (10-11%).<sup>4</sup> By race, 75-79% of AHCCCS children reported as Caucasian/White, 3% as Black, 2% as Asian/Pacific Islander, and 3-4% as Native American. Of these, 74-75% of AHCCCS children were Hispanic or Latino, and 5% were affiliated with a tribal community.<sup>5</sup> Of the regional AHCCCS children, 82-91% were enrolled in two health plans: Steward Health Choice AZ (62-63%) and UnitedHealthcare (20-29%). Most annual health claims were submitted by physicians (29-32%), pharmacies (15-16%), behavioral health outpatient clinics (9-12%), and hospitals (10%).

## **Health Care Workforce**

The supply of physicians in the United States is tracked by the Association of American Medical Colleges biennially. Arizona had 160 hospitals individually licensed by the state which were subtyped as children, critical access, long term, short term, psychiatric, rehabilitation, transplant and non-participating. The La Paz/Mohave Region had four short-term hospitals, six Federally Qualified Health Center (FQHC) sites<sup>6</sup>, six rural health clinic sites<sup>7</sup>, ambulatory surgical centers, medical groups, outpatient treatment centers, and other medical facilities available in the region. The rate of available primary care physicians in the region was 10-14 primary care physicians per 1,000 AHCCCS children compared to the statewide rate of 23-24 per 1,000 AHCCCS children. For primary care physicians accepting AHCCCS patients, the regional rate was 8-10 physicians per 1,000 AHCCCS children. For dentists accepting AHCCCS patients, the regional rate was 3 dentists per 1,000 AHCCCS children compared to 16-17 dentists per 1,000 AHCCCS children statewide.

We compared the distance that regional and statewide AHCCCS children needed to travel to the nearest provider type to assist in determining whether the population in the region may have access to care issues based on travel distance. To visit the nearest primary care physician or pharmacy or dentist, 21-29% of AHCCCS children in the region traveled up to one mile for services and another 50-57% traveled one to five miles compared to 56-64% of AHCCCS children statewide who traveled up to one mile and another 26-35% traveled one to five miles. The nearest hospital was one mile away for 10% of regional AHCCCS children, one to five miles away for 66% and more than 10 miles away for 17-18% compared to 11-12% of statewide AHCCCS children who traveled one mile or less, 69% who traveled one to five miles and 10% who traveled 10 miles or more. To visit the nearest behavioral health provider, 38% of regional AHCCCS

<sup>&</sup>lt;sup>4</sup> The La Paz/Mohave Remainder subregion had <6 AHCCCS children; subsequently, this subregion is excluded from reporting after this section.

<sup>&</sup>lt;sup>5</sup> Tribal affiliation refers to whether an individual is a member of a federally recognized Arizona tribe and is displayed as a flag (Yes/No) in this report. This information is captured during enrollment in AHCCCS.

<sup>&</sup>lt;sup>6</sup> The FQHC sites are Creek Valley Health Clinic in Colorado City, North Country Healthcare (two in Kingman, one in Bullhead City and one in Lake Havasu City) and River Cities Community Clinic in Bullhead City.

<sup>&</sup>lt;sup>7</sup> The Rural Health Clinics are La Paz Regional Hospital (one in Quartzsite and one in Parker), Parker Medical Center, Parker Walk-in Clinic, San Luis Walk-in Clinic in Parker and Tri-Valley Medical Center in Salome.

children traveled one mile and 43% traveled one to five miles; for statewide AHCCCS children, 62-65% traveled one mile to the nearest behavioral health provider and 27-30% traveled one to five miles.

## **Primary Care and Well-Child Visits**

Access to primary care is important for the health and well-being of children. Primary care practitioners (PCPs) provide appropriate screenings, treatment and preventive services. When children regularly visit a PCP, they are less likely to visit the emergency department for non-urgent care. Well-child visits are PCP visits scheduled at designated age intervals where a child's growth and development are measured and tracked according to national guidelines. PCPs examine a child holistically for physical, mental, emotional and social/environmental health during a well-child visit.

Regionally, 81-83% of AHCCCS children had at least one PCP visit compared to 85-86% of AHCCCS children statewide and 86-87% of Medicaid children nationally. The regional rates were below the AHCCCS Minimum Performance Standard (MPS)<sup>8</sup> of 84%. The subregions who met or exceeded the MPS for annual PCP visits included Bullhead City (all years), Dolan Springs (2019), Fort Mohave (2019) and Lake Havasu City (2017 and 2019). Regional AHCCCS children who had annual PCP visits were more likely to be ages 1-2 (88-90%) than ages 3-5 (76-79%) and Hispanic and Latino (84%) than Non-Hispanic or Latino (80-82%).

Regionally, 46-53% of AHCCCS children birth to 15 months had six or more well-child visits compared to 53-60% of AHCCCS children statewide and 63-66% of Medicaid children nationally. Approximately 95% of all regional AHCCCS children had at least one well-child visit. The subregions who met or exceeded the AHCCCS statewide rates for six or more well-child visits for AHCCCS children birth to 15 months were Bullhead City (2017 and 2018) and Lake Havasu City (2017 and 2019). Regional AHCCCS children birth to 15 months who had six or more well-child visits were more likely to be Hispanic or Latino (48-61%) than Non-Hispanic or Latino (46-50%) and Asian/Pacific Islander in 2017 and 2019 (58-85%) than other races. For AHCCCS children ages 3-5, 51-56% of regional children had an annual well-child visit compared to 62-65% of statewide children and 72-74% of Medicaid children ages 3-6 nationally. No subregion met the AHCCCS statewide rates for this indicator. Regional AHCCCS children ages 3-5 receiving an annual well-child visit were more likely to be Hispanic or Latino (55-58%) than Non-Hispanic or Latino (49-55%).

## **Screening for Lead Poisoning**

Lead poisoning is a silent killer because often there are no symptoms. Exposure to lead can cause irreversible damage to the brain and other vital organs in children, as well as intellectual and behavioral deficits. To detect abnormal blood lead levels in children, screenings are conducted via a blood lead test. According to the Arizona Department of Health Services (ADHS), children who live in areas designated as

Contractors. AHCCCS-reported rates are the official rates used to determine Contractor compliance with performance requirements. If a Contractor does not achieve the MPS, they will be required to submit a corrective action plan and may be subject to sanctions for each deficient measure.

<sup>8</sup> Minimum Performance Standard (MPS) is the minimal expected level of performance by AHCCCS

high-risk for lead poisoning should receive a blood lead test at 12 and 24 months of age, and older children who have not been previously tested should receive the blood lead test.<sup>9</sup>

ADHS reported 61,391 children under age six (14% of children under age 5) were screened in 2019, and 40,773 (66%) of those children lived in high-risk areas. Of the children living in high-risk areas, 29% were screened at 12 months of age, and 19% were screened at 24 months of age. Only 10% of children were screened at both intervals. For AHCCCS children being screened for lead poisoning one or more times by their second birthday in, the regional rates increased from 23% in 2017 to 35% in 2019 compared to AHCCCS statewide rates which increased from 32% in 2017 to 35% in 2019. Two subregions met or exceeded the AHCCCS statewide rates for one or more blood lead screenings for lead poisoning by the second birthday: Fort Mohave (2018 and 2019) and Lake Havasu City (2017). Hispanic or Latino AHCCCS children (28-36%) were more likely to be screened for lead poisoning by their second birthday than Non-Hispanic or Latino AHCCCS children (21-35%).

## Weight Assessment and Counseling

Childhood obesity has both short-term and long-term effects, so it is important for PCPs to monitor weight problems in children and provide guidance for maintaining a healthy weight and lifestyle. The prevalence of obesity among children aged 2–5 years in 2015-2016 was 14% according to a national survey. For this report, we focused on AHCCCS children ages 3-5.

The regional rates for weight assessment  $^{10}$  and counseling showed AHCCCS children in the La Paz/Mohave Region were assessed for weight at 6-7% compared to AHCCCS children statewide who were assessed at rates of 9-19%. Rates for nutrition counseling were 1-2% at the regional level versus 4-5% at the state level for AHCCCS children. Physical activity assessments in the region were suppressed, and the state was assessed at rates of <1-1%.  $^{11}$ 

#### **Developmental Screening and Delay**

During early childhood, children grow and develop at a rapid pace physically and cognitively. Although children develop skills at different times, there are guidelines that define the period when an average child should meet certain developmental milestones. National pediatric guidelines recommend developmental screenings during well-child visits for all children ages 9 months, 18 months, 2 years and

zip codes.

<sup>&</sup>lt;sup>9</sup> See <a href="https://www.azdhs.gov/preparedness/epidemiology-disease-control/lead-poisoning/index.php#high-risk-zip-codes-home">https://www.azdhs.gov/preparedness/epidemiology-disease-control/lead-poisoning/index.php#high-risk-zip-codes-home</a> for an interactive neighborhood map to identify high-risk

<sup>&</sup>lt;sup>10</sup> Under HEDIS, the rates for weight assessment are an evaluation of whether Body mass index (BMI) percentile is accessed and does not determine the absolute BMI value. The diagnosis codes for pediatric BMI included: Z68.51 (< 5th percentile for age), Z68.52 (5th percentile to < 85th percentile for age), Z68.53 (85th percentile to < 95th percentile for age) and Z68.54 (≥ 95th percentile for age).

<sup>&</sup>lt;sup>11</sup> Physical Activity Counseling includes sports physicals which are not provided to children in the early childhood age group.

2.5 years. Developmental delay occurs when a child does not demonstrate mastery of developmental milestones. Developmental delays have been found to occur in 10-15% of preschool children nationwide.

Rates of developmental screenings in AHCCCS children birth to age 5 increased over the report period although still well below the AHCCCS median rates. At the regional level, rates increased 10-16% compared to statewide AHCCCS rates of 10-14%. <sup>12</sup> The subregions with the highest rates of developmental screenings for AHCCCS children were Dolan Springs (2017 and 2019), Kingman (all years) and Lake Havasu City (2017 and 2019). Regional AHCCCS children who received developmental screenings were more likely to be ages 1-2 (28-42%) than age 0 (5-9%) or ages 3-5 (1-4%), and Asian/Pacific Islander in 2017 and 2018 (55-58%) than other races.

Rates of diagnosing developmental delay in AHCCCS children were 2-4% at the regional level compared to 3-5% at the state level for AHCCCS children. Regional AHCCCS children who were diagnosed with developmental delay were more likely to be male (3-4%) than female (2-4%), and ages 3-5 (3-5%) than age 0 (1%) or ages 1-2 (2-4%). Of those AHCCCS children who were diagnosed with developmental delay, 42-50% of regional AHCCCS children received behavioral health services compared to 47-58% of AHCCCS children statewide. The subregions where AHCCCS children who were diagnosed with developmental delay were most likely to receive behavioral health services included Bullhead City (2017 and 2018) and Dolan Springs (2018). Regional AHCCCS children who were diagnosed with developmental delay and received behavioral health services were more likely to be male (50-59%) than female (30-39%), ages 3-5 (45-65%) than ages 1-2 (31-40%) and affiliated with a tribe (60-75%) than not affiliated (41-50%).

#### **Behavioral Health**

The social-emotional development and adaptive functioning of a young child is as important as their physical health. Negative early childhood events can lead to behavioral and physical health problems in adulthood if behavioral health intervention services are not provided at the infant and toddler stages. For young children, behavioral health services <sup>13</sup> would likely include day programs, crisis services, rehabilitation services, health promotion, mental health counseling, psychiatric and psychologist services, and various support services.

Eleven to sixteen percent of regional and statewide AHCCCS children received behavioral health services. The subregions where AHCCCS children were more likely to receive behavioral health services included Colorado City (2019), Kingman (2019) and Quartzsite (2017). Male AHCCCS children in the region were more likely to receive behavioral health services (13-19%) then females (8-13%).

\_

<sup>&</sup>lt;sup>12</sup> Due to the limited capture of developmental screenings in claims data alone, these rates should be interpreted with caution.

<sup>&</sup>lt;sup>13</sup> Visit <a href="https://www.azahcccs.gov/Members/AlreadyCovered/coveredservices.html">https://www.azahcccs.gov/Members/AlreadyCovered/coveredservices.html</a> for more details on AHCCCS behavioral health services.

#### Vision<sup>14</sup>

Visual impairment affects a child's development, performance, and quality of life. Fortunately, most vision problems are successfully treated when detected early through regular visits to PCPs, and well-child visits should include a vision screening. It has been estimated that 20% of preschool children in the United States have eye or vision problems. Arizona's Eyes on Learning Vision Coalition recommends a vision screening beginning at age one. Children ages 3-5 should have at least one vision screening by a PCP or trained screener, and annual screenings should be provided to children in kindergarten through fourth grade.

In the La Paz/Mohave Region, 45-48% of AHCCCS children received an annual vision screening or well-child visit compared to 43-47% of AHCCCS children statewide. Regional AHCCCS children who received an annual vision screening or well-child visit were more likely to be ages 1-2 (69-74%) than ages 3-5 (51-54%) and Hispanic or Latino (49-53%) than Non-Hispanic or Latino (44-47%). The subregions where AHCCCS children were more likely to receive an annual vision screening or well-child visit included Bullhead City (all years), Dolan Springs (all years), Fort Mohave (all years), Kingman (all years) and Lake Havasu City (2017).

Eye exams were conducted much less frequently, ranging 5-7% annually at the regional level and 4-5% at the state level for AHCCCS children. Regional AHCCCS children ages 3-5 (10-11%) were more likely to receive an eye exam than ages 1-2 (2-4%). Follow-up eye exams were conducted on AHCCCS children in the region and statewide at rates of 4-5%. AHCCCS children with visually significant eye conditions received treatment at rates of 53-67% regionally compared to 54-60% statewide.

## Hearing<sup>14</sup>

Most children begin hearing sounds at birth and learn to speak over time by imitating the sounds around them. However, it is reported that around two or three out of every 1,000 children are born deaf or hard-of-hearing in the United States, and more lose their hearing later in childhood. For children diagnosed with hearing loss, early detection, intervention and treatment would provide each child with the opportunity to develop better language and communication skills. Arizona strives to screen all infants before one month of age. Infants who do not pass the initial hearing screen and a rescreening, should be evaluated further to confirm or diagnose hearing loss before 3 months of age. Infants diagnosed with permanent hearing loss should receive intervention services before 6 months of age.

Around 99% (82,035) of all Arizona infants received a newborn hearing screening in 2017, which was slightly higher than the national rate of 98%. Less than 1% of all Arizona infants were diagnosed with permanent hearing loss, and of those, 42% were diagnosed before three months of age. Nationally, 10%

<sup>14</sup> ibid

\_

<sup>&</sup>lt;sup>14</sup> Per the AHCCCS Medical Policy Manual, AHCCCS children should receive hearing and vision screenings during their well-child visits according to the periodicity schedule. Claims data does not specify each service provided during a well-child visit; thus, we cannot verify whether these screenings were provided according to the schedule. The rates in this report should be interpreted with caution.

of infants were diagnosed with permanent hearing loss, and of those, approximately 74% were diagnosed before three months of age. Around 42% of Arizona infants were diagnosed with hearing loss before three months of age compared to approximately 74% nationally. Additional audiology services were provided to 4-9% of AHCCCS children under age one in the La Paz/Mohave Region compared to 9-12% of AHCCCS children statewide. Regional AHCCCS children under age one were more likely to receive additional audiology services in Fort Mohave in all years.

Hearing screenings were provided to 10-11% of AHCCCS children ages 1-5 in the region compared to 20-28% of AHCCCS children statewide. Regional AHCCCS children ages 1-5 were more likely to receive hearing screenings included Dolan Springs (2017) and Lake Havasu City (2018 and 2019). Male regional AHCCCS children ages 1-5 were more likely to receive hearing screenings (12-14%) than females (8-9%). The provision of additional audiology services ranged 72-88% of AHCCCS children in the region while statewide AHCCCS children's rates decreased from 68% to 57% over the same period. The subregions where AHCCCS children ages 1-5 were more likely to receive additional audiology services included Dolan Springs (2017 and 2018), Fort Mohave (2019), Kingman (2017 and 2019) and Lake Havasu City (all years).

#### **Oral Health**

Oral health is a key indicator of overall health, well-being and quality of life. Access to dental care is necessary to maintain good oral health. Two preventative care dental visits are recommended annually for children. For young children, the application of fluoride varnish to primary and permanent teeth every six months is recommended to prevent cavities.

In La Paz/Mohave, 51-56% of AHCCCS children had at least one annual dental visit compared to 51-53% of AHCCCS children statewide. Neither the region nor the state met the AHCCCS Minimum Performance Standard (MPS) of 60% for annual dental visits for ages 2-20. Colorado City and Kingman subregions met or exceeded the AHCCCS MPS in 2019. Regional AHCCCS children with at least one annual dental visit were more likely to be ages 3-5 (61-65%) than ages 1-2 (36-41%) and Hispanic or Latino (54-58%) than Non-Hispanic or Latino (51-55%).

Two preventative care dental visits are recommended annually for children. In the region, 22-27% of AHCCCS children received the biannual preventative care dental visit compared to 18-20% of AHCCCS children statewide; and over half of regional AHCCCS children had at least one preventative care dental visit per year. Fluoride varnish was applied to 45-52% of AHCCCS children in the region compared to 47-49% of AHCCCS children statewide. The subregions that met or exceeded the AHCCCS statewide rates for AHCCCS children having received a fluoride varnish application included Bullhead City (all years), Dolan Springs (all years), Fort Mohave (all years) and Kingman (all years). Regional AHCCCS children who had a fluoride varnish application were more likely to be ages 3-5 (55-60%) than ages 1-2 (30-38%), Hispanic or Latino (49-55%) than Non-Hispanic or Latino (44-51%) and Asian/Pacific Islander (50-57%).

## **Maternal Prenatal and Postpartum Care**

The health of women before pregnancy and after delivery significantly impacts their health and the health of their babies. Thus, it is important to focus on women's prenatal and postpartum care. Prenatal care involves regular visits to a health care provider to monitor the mother's health and health of the developing fetus. Women should have at least one prenatal visit in the first trimester of pregnancy. The

period of up to 60 days following childbirth is called the postpartum period for AHCCCS women. Preexisting health conditions, social determinants, and newly developed conditions contribute to maternal morbidity and mortality during this period so at least one postpartum visit is recommended.

In La Paz/Mohave, 94-95% of pregnant women began prenatal care in the first trimester compared to 84-86% of AHCCCS women statewide, which were both above the Healthy People 2030 target rate of 81%<sup>15</sup>. For postpartum care, 96-97% of regional AHCCCS women had at least one postpartum visit compared to 88-89% of AHCCCS women statewide and 64-75% of Medicaid women nationally. Most subregions met or exceeded the Healthy People 2030 target rate for timely prenatal care in all years and postpartum visits. The demographic breakdowns for AHCCCS women's prenatal and postpartum care showed high rates, ranging 90-100%.

#### **Health Plan Performance**

This section provided a selection of health indicators to compare results among AHCCCS children enrolled in each of the AHCCCS health plans available in the region. Six regional health plans met or exceeded the AHCCCS statewide aggregate performance and MPS for PCP visits for ages 1-5: Care 1st (2017), Children's Rehabilitative Services (2017 and 2018), Comprehensive Medical and Dental Program (CMDP) (2018 and 2019), DES DD (all years), and Mercy Care Plan (all years) and UnitedHealthcare (2019). Regional performance for at least one well-child visit in the first 15 months was high in at least one year for all reporting health plans. Three health plans met or exceeded the AHCCCS statewide aggregate performance and MPS for annual well-child visits for ages 3-5: CMDP (2018), DES DD (2019) and Mercy Care Plan (2018). CMDP exceeded the MPS for preventative care dental visits for ages 1-5 in 2017 and 2018.<sup>16</sup>

#### Conclusion

From 2017 to 2019, the La Paz/Mohave Region showed strong performance on the following AHCCCS children's and women's health indicators: newborn hearing screenings, immunizations (DTaP, Hepatitis A, Combo 3), vision screenings, and prenatal and postpartum care. These achievements contributed to good health outcomes throughout the region. The areas where needs were identified for AHCCCS women and children included access to care, the supply of health care professionals, PCP visits, developmental screenings, well-child visits, lead poisoning screenings, hearing screenings for ages 1-5 and oral health. The information in this report can be combined with other available information to create a more comprehensive view of young children and women in the region for regional council planning.

<sup>&</sup>lt;sup>15</sup> See https://health.gov/healthypeople/objectives-and-data/browse-objectives/pregnancy-andchildbirth/increase-proportion-pregnant-women-who-receive-early-and-adequate-prenatal-care-mich-08 for the Healthy People 2030 Prenatal Care Objective.

<sup>&</sup>lt;sup>16</sup> The AHCCCS statewide indicator for preventative care dental visits includes ages 2-20 which incorporates a significantly larger number of AHCCCS children than our reporting on ages birth to 5, so the rates should be compared with caution.

## **INTRODUCTION**

## THE IMPORTANCE OF EARLY CHILDHOOD HEALTH

Under the direction of First Things First (FTF), the Arizona State University Center for Health Information & Research (CHiR) conducted a regional analysis of children from birth to age 5 to explore the health assets and needs in the FTF La Paz/Mohave Region. This report provides detailed health utilization and access to services for children birth through 5, along with prenatal and postpartum women, who were enrolled in the Arizona Health Care Cost Containment System (AHCCCS). Additional information is provided via medical board licensing data to further describe access to medical professionals and services contracted under AHCCCS. The goal is for the FTF Regional Partnership Councils to utilize the findings in this report when conducting regional planning discussions, as an additional resource and tool to the Regional Needs and Assets base report.

#### **DEFINITIONS**

## Access to Care

This term refers to making health care services readily available when needed and removing all barriers.

## Age Groups

- Age is defined as the age of the patient on Dec. 31 of each given year. The age for a patient is constant through the report year. Special age breakdowns are listed for certain indicators that have an associated milestone.
- Infants: less than 1 year of age
- Toddlers: greater than or equal to 1 year of age to less than 3 years of age
- Preschooler: greater than or equal to 3 years of age to less than 6 years of age

#### Assets

An asset is a finding where young children or women appear to be faring well regarding utilization of or access to health care.

#### Behavioral Health

To determine whether children are receiving behavioral health services, we used the following definition: category of service on claim equals mental health services (category of service = 47) or primary diagnosis is a behavioral health diagnosis as listed in the AHCCCS Behavioral Health Services Matrix <a href="https://www.azahcccs.gov/PlansProviders/MedicalCodingResources.html">https://www.azahcccs.gov/PlansProviders/MedicalCodingResources.html</a>.

#### Children

Unless noted otherwise, all references to children denote children, ages birth to 5, who are AHCCCS members.

#### CMS Median

The Centers for Medicare & Medicaid Services (CMS) annually collects and reports state performance rates on a standardized set of care quality measures for Medicaid and Children's Health Insurance Program beneficiaries, called the Child and Adult Core Set. The CMS Median is the average performance among reporting states for each measure.

#### Habilitation

Training in independent living skills or special developmental skills, sensory-motor development, orientation and mobility and behavior intervention.

## <u>Healthcare Effectiveness Data and Information Set</u>

The Healthcare Effectiveness Data and Information Set (HEDIS) is a tool produced by the National Committee for Quality Assurance (NCQA) that is used by most U.S. health plans to measure performance and quality in health care. HEDIS® contains over 90 measures under six domains of care: effectiveness of care, access/availability of care, experience of care, utilization and risk adjusted utilization, health plan descriptive information, and measures reported using electronic clinical data systems. The national committee collects HEDIS® survey results from health plans and Preferred Provider Organizations through the Healthcare Organization Questionnaire and collects non-survey data through the Interactive Data Submission System. HEDIS measures are specifically defined to make comparisons among health plans. The measurement set is reviewed annually. CHiR uses the AHCCCS claims within HEDIS. HEDIS measures have complicated numerator and denominator calculations, and therefore, are expressed and interpreted as rates.

#### **Health Plans**

Health plan categories include acute care, Children's Rehabilitative Services, Comprehensive Medical and Dental Program, Developmental Disability/Department of Economic Security, Long Term Care, and Fee-For-Service American Indian health plans.

## Minimum Performance Standard

Minimum Performance Standard (MPS) is the minimal expected level of performance by AHCCCS Contractors. AHCCCS-reported rates are the official rates used to determine Contractor compliance with performance requirements. If a Contractor does not achieve the MPS, they will be required to submit a corrective action plan and may be subject to sanctions for each deficient measure.

## Needs

A need is an area where it appears that access or utilization of health care is low.

#### <u>Outpatient Treatment Center</u>

Under Arizona Administrative Code R9-10.101.39, an "outpatient treatment center" means a healthcare institution class without inpatient beds that provides medical services for the diagnosis and treatment of patients. These are generally small private or public health facilities whose services may include primary care, wellness and prevention, diagnostics, treatment and minor surgeries (i.e., Urgent Care Centers, Dialysis Centers, Primary Care Centers and Imaging Centers).

#### Postpartum Period

The AHCCCS postpartum period begins the day the pregnancy terminates and continues for 60 days following pregnancy termination.

## **Primary Care Physician Specialties**

Physicians included in the primary care specialty include Family Practitioner, General Practitioner, Internal Medicine and Pediatrician.

## Race/Ethnicity

Race and ethnicity are grouped and reported in the following manner.

- Race
  - Asian/Pacific Islander
  - o Black
  - Caucasian/White
  - o Native American
  - Other/Unknown

- Ethnicity
  - Hispanic or Latino
  - Not Hispanic or Latino
  - o Unknown

Up until 2017, AHCCCS only collected one race/ethnicity variable and used the Hispanic value to denote Hispanic or Latino origin. As of 2017, AHCCCS began collecting race and ethnicity as separate variables. Hispanic is retained as a race variable, but AHCCCS is phasing out its use; therefore, the decrease in the use of Hispanic in the race variable correlates to the increase in reporting of Unknown in the race variable. Ethnicity is reported separately beginning in 2018 and notes on its use in this report are below.

- Individuals who reported "Not Hispanic, Latino, Spanish" are not Hispanic or Latino origin.
- To denote those of Hispanic or Latino origin, we combine Mexican, Mexican American, Chicano, Puerto Rican, Cuban, Other Hispanic/Latino Origin, and Hispanic or Latino Unknown.
- All individuals who reported a race/ethnicity of Hispanic prior to 2017 were assigned a race of other/unknown and an ethnicity of Hispanic or Latino origin
- Ethnicity Unspecified refers to individuals who did not answer or were not provided the opportunity to give this information.
- Ethnicity Unknown means the individual chose to be unknown. AHCCCS started phasing out this category in October 2018.
- Data on multiracial individuals is not collected.

## **Tribal Affiliation**

Tribal affiliation refers to whether an individual is a member of a federally recognized Arizona tribe and is displayed as a flag (Yes/No) in this report. This information is captured during enrollment in AHCCCS. This is based on AHCCCS-stated affiliation, not residential location.

## Well-Child Visits

Children enrolled in AHCCCS receive well-child visits under the Early and Periodic Screening, Diagnostic and Treatment (EPSDT) Program. The EPSDT program provides comprehensive treatment and preventive

health care services for children under age 21. The services include dental, physical, behavioral health, developmental, vision, hearing, screenings and other specialty services. EPSDT visits are all-inclusive, meaning one payment is made for all services rendered during the visit. Only certain services are billed separately when conducted by qualified health care providers, and those are: nutritional assessments, developmental screenings, immunizations, fluoride varnish and ocular photo screening.

#### Women

Unless noted otherwise, all references to women denote women who were AHCCCS members.

#### **APPROACH**

CHiR and representatives from the FTF Regions, Programs, and Evaluation teams determined priority indicators for this report. FTF provided the regional and subregional boundaries. The La Paz/Mohave Region consisted of 11 subregions: Bullhead City, Colorado City, Dolan Springs, Fort Mohave, Kingman, Lake Havasu City, La Paz/Mohave Remainder, Littlefield, Parker, Quartzsite and Salome. The main data source was claims data from the Arizona Health Care Cost Containment System (AHCCCS), Arizona's Medicaid agency; therefore, the results presented in this report were for children and mothers who were enrolled in AHCCCS from 2017 to 2019. This population was denoted as AHCCCS children or AHCCCS women.

AHCCCS children's health was measured in the following categories: primary care, well-child visits, health care workforce, screening for lead poisoning, weight assessment and counseling, developmental screening and delay, behavioral health, vision, hearing, oral health, immunizations, maternal prenatal and postpartum care, and health plan performance.

Many of the reported indicators were from the Healthcare Effectiveness Data and Information Set (HEDIS). HEDIS is a performance improvement tool whereby health plans, health care organizations and government agencies submit data on specific health measures. HEDIS used the collected data to calculate national performance statistics and benchmarks and set standards for measures. HEDIS specifications were applied to the AHCCCS population for each region. The denominators were listed within each indicator and are available on the National Committee for Quality Assurance website at <a href="https://www.ncqa.org/hedis/">https://www.ncqa.org/hedis/</a>. Inclusion generally required a child to have continuous enrollment for the reporting year with no more than one gap smaller than 45 days allowed. Some indicators also required enrollment in a period preceding the reporting year.

Non-HEDIS denominators were derived from the children who met the AHCCCS inclusion criteria for the region. The AHCCCS inclusion criteria were children ages 0-5 (0 <= age < 6) who were enrolled in AHCCCS in 2017, 2018, or 2019 and residing in Arizona regions defined by First Things First. Health claims were for paid services in 2017, 2018, or 2019. Additional AHCCCS enrollment requirements were indicator-based. The complete population of children covered by AHCCCS were not included due to the limitations on AHCCCS enrollment gaps which were not met by all children.

For the distance analysis that was reported in the health care workforce section, all AHCCCS-enrolled children were assigned coordinates on a map related to their residential address on file. Health providers

were also assigned coordinates from their address on file or public address, if available. Each child's address was analyzed to determine the distance in miles to the closest provider for each provider type. The children were then grouped into distance ranges as percentages. The region and state percentages were listed side-by-side to compare totals and determine if the population in the region may have access to care issues due to the distance required to travel for health services.

Data used in this report covered all AHCCCS members in Arizona, including members living in FTF tribal regions and subregions. Report creations for tribal regions and subregions was carried out with specific approval from the tribe. As required by a Resolution previously established between FTF and the Fort Mojave Indian Tribe, which is located within the FTF La Paz/Mohave Region, FTF notified the Fort Mojave Indian Tribe Region in writing that work would begin on this report, and the work was approved by the region. For those tribal regions who did not give approval, data was included only in aggregate totals for Arizona. In the case of a tribal subregion, only aggregate totals for Arizona and the region were included.

#### REPORTING

There were 13 health topics discussed in this report. Each section began with context on the importance of the health topic before discussing the results from the AHCCCS claims data.

The AHCCCS results were presented at the regional level with state and national benchmarks provided for comparison, where available. When possible, the results are grouped by 1) indicators that met or were above the state average or national HEDIS standards and 2) indicators that did not meet or were below the state average or national HEDIS standards. Other notable findings were also presented that do not have comparison data. Most results were presented as percentages for standardization purposes and ease of comparison with benchmarks. The terms rate and percent were used interchangeably.

After reporting the general regional demographics, the results were displayed by gender, age group, race, ethnicity, tribal affiliation, provider type, and/or health plan when the data was available and within the data suppression guidelines stated below. Each section contained maps to display the results at the subregional level. The maps had a color gradient which compared the performance among the subregions for each indicator. A darker color denoted a higher percentage of individuals in the subregion who were included in the indicator. Percentages over 1% were rounded to the nearest whole number. Percentages less than 1% were denoted as "<1%".

A brief conclusion summarized how well the region was doing with regards to access and utilization of health care services and provided areas where the regional councils may want to focus during their regional planning conversations.

The Executive Summary was designed to provide the main findings and takeaways for the report. A definitions section explained the lesser-known terms. The data sources were detailed in the Appendix which follows the references. The report was hyperlinked for ease of navigating from the Table of Contents and the text to the associated topics, figures and tables.

To protect the confidentiality of program participants, the First Things First Data Dissemination and Suppression Guidelines preclude our reporting data related to health or developmental delay if the count

is less than six. Throughout this report, information which was not available because of suppression guidelines will be indicated by entries of "<6" for counts or "DS" (data suppressed) for percentages. Data were sometimes not available for particular regions, either because a particular program did not operate in the region or because data were only available at a higher level (i.e., county, state, etc.). Cases where data were not available will be indicated by an entry of "N/A."


#### **DATA LIMITATIONS**

Most of the results in this report used AHCCCS claims and encounter data. While being limited to the population of children enrolled in Arizona Medicaid, this data source was also subject to coding errors and missing data for some indicators.

To best capture the full picture of childhood immunizations, a combination of data from claims, electronic health records, paper medical records and registry data was needed. AHCCCS used a combination of AHCCCS claims, Arizona State Immunization Information System (ASIIS) registry data and medical record data from its contractors to measure immunization rates internally. For this report, we used only AHCCCS claims as we did not have access to the other data sources. Since the AHCCCS claims data only included a subset of the immunizations of Arizona's children, our results showed substantially lower immunization rates than AHCCCS officially reports.

Per the AHCCCS Medical Policy Manual, AHCCCS children should receive hearing and vision screenings during their well-child visits according to the periodicity schedule. Claims data does not specify each service provided during a well-child visit; thus, we cannot verify whether these screenings were provided according to the schedule. The rates in this report should be interpreted with caution.

# LA PAZ/MOHAVE REGIONAL RESULTS



#### POPULATION AND DEMOGRAPHICS OF CHILDREN ENROLLED IN AHCCCS

AHCCCS exists to make care affordable to the individuals and families it enrolls, including the approximately 235,000 children birth to age 5 who were enrolled in AHCCCS from 2017 to 2019.

In the La Paz/Mohave Region, there were 6,970 children from birth to age 5 enrolled in AHCCCS in 2017, 6,701 children enrolled in 2018 and 6,606 enrolled in 2019.

(AHCCCS Claims Data, 2021)

Of the La Paz/Mohave children enrolled in AHCCCS from 2017 to 2019, male AHCCCS children outnumbered females by 3-4% (Table 1). In Table 2, there were 7% more infants and toddlers than preschoolers in 2017, with the difference dropping to 5% more in 2018 and 3% more in 2019.

Approximately 87% of the AHCCCS children in the region lived in four subregions: Kingman (32-33%), Bullhead City (23%), Lake Havasu City (19-20%) and Fort Mohave (10-11%) (Figure 1).<sup>17</sup> By race, 75-79% of AHCCCS children reported as Caucasian/White, 3% as Black, 2% as Asian/Pacific Islander, and 3-4% as Native American (Figure 2). Of these, 74-75% of AHCCCS children were Hispanic or Latino (Figure 3), and 5% were affiliated with a tribal community (Figure 4).<sup>18</sup> In Figure 5, 82-91% of AHCCCS children in the region were enrolled in two health plans: Steward Health Choice AZ (62-63%) and UnitedHealthcare (20-29%). Most annual health claims were submitted by physicians (29-32%), pharmacies (15-16%), behavioral health outpatient clinics (9-12%), and hospitals (10%).

Table 1. Number of AHCCCS Children Birth to Age 5 by Year and Sex

Year	Female	Male	Total of AHCCCS-Enrolled Children
2017	3,382	3,588	6,970
2018	3,217	3,484	6,701
2019	3,191	3,415	6,606

Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

Table 2. Number of AHCCCS Children Birth to Age 5 by Year and Age Group

Year	Infant (under 1) Toddler (1-2)		Preschooler (3-5)
2017	1,464	2,251	3,255
2018	1,483	2,042	3,176
2019	1,414	1,989	3,203

Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

<sup>17</sup> The La Paz/Mohave Remainder subregion had <6 AHCCCS children; subsequently, this subregion is excluded from reporting after this section.

<sup>&</sup>lt;sup>18</sup> Tribal affiliation refers to whether an individual is a member of a federally recognized Arizona tribe and is displayed as a flag (Yes/No) in this report. This information is captured during enrollment in AHCCCS.




Figure 1. Number of AHCCCS Children Birth to Age 5 by Year and Subregion

Asian/Pacific Islander 117 Black 240 Caucasian/White 5,489 Native American 258 Other/Unknown 866 2018 Asian/Pacific Islander 112 Black 232 Caucasian/White 5,215 Native American 239 Other/Unknown 903 2019 Asian/Pacific Islander 114 Black 230 Caucasian/White 4,924 Native American 224 Other/Unknown 1,114 0K 1K 2K 3K 4K 5K 6K Count of Children

Figure 2. Number of AHCCCS Children Birth to Age 5 by Year and Race




Figure 3. Number of AHCCCS Children Birth to Age 5 by Year and Ethnicity

2017 2018 2019 7K 6,612 6,358 6,260 Count of Children 6K 5K 4K 3K 2K 1K 358 343 346 No Yes No Yes No Yes

Figure 4. Number of AHCCCS Children Birth to Age 5 by Year and Tribal Affiliation

Figure 5. Number of AHCCCS Children by Year and Health Plan




Table 3. Number of AHCCCS Claims by Provider Type (Billing Entity), 2017-2019

	20	17	20	18	2019		
Provider Type	Claims Count	Percent of Total	Claims Count	Percent of Total	Claims Count	Percent of Total	
Behavioral Health Outpatient Clinic	11,125	12%	7,920	9%	8,802	9%	
Dentist	5,146	6%	4,986	6%	5,404	6%	
Durable Medical Equipment Supplier	717	1%	825	1%	1,011	1%	
Federally Qualified Health Center (FQHC)	664	1%	1,114	1%	1,370	1%	
Habilitation Provider*	2,750	3%	2,437	3%	2,595	3%	
Home Health Agency	241	<1%	452	1%	902	1%	
Hospital	8,849	10%	8,252	10%	9,053	10%	
Integrated Clinics**	916	1%	954	1%	802	1%	
Laboratory	1,608	2%	1,750	2%	1,623	2%	
Non-Emergency Transportation Providers	475	1%	442	1%	491	1%	
Occupational Therapist	1,799	2%	2,032	2%	1,815	2%	
Pharmacy	14,737	16%	13,469	16%	14,054	15%	
Physical Therapist	837	1%	968	1%	1,197	1%	
Physician – MD/DO	29,202	32%	25,586	29%	27,826	30%	
Physician Assistant	2,297	3%	2,886	3%	2,088	2%	
Registered Nurse Practitioner	5,016	6%	6,583	8%	7,281	8%	
Speech Language Pathology	1,058	1%	2,722	3%	2,394	3%	
Speech/Hearing Therapist	1,664	2%	2,083	2%	2,800	3%	
Other	1,434	2%	1,328	2%	1,535	2%	

Notes: \* Habilitation is training in independent living skills or special developmental skills, sensory-motor development, orientation and mobility and behavior intervention.

<sup>\*\*</sup> An Integrated Clinic is a provider licensed by the Arizona Department of Health Services as an Outpatient Treatment Center which provides both behavioral health services and physical health services.

#### **HEALTH CARE WORKFORCE**

Currently, Arizona has 160 hospitals individually licensed by the state which are subtyped as children, critical access, long term, short term, psychiatric, rehabilitation, transplant and non-participating (Arizona Department of Health Services, 2021). The La Paz/Mohave Region had four short-term hospitals, six Federally Qualified Health Center (FQHC) sites<sup>19</sup>, six rural health clinic sites<sup>20</sup>, ambulatory surgical centers, medical groups, outpatient treatment centers, and other medical facilities available in the region (Arizona Department of Health Services, 2021).

The rate of available primary care physicians in the region was 10-14 primary care physicians per 1,000 AHCCCS children (Table 5) compared to the statewide rate of 23-24 per 1,000 AHCCCS children (Table 4). For primary care physicians accepting AHCCCS patients, the regional rate was 8-10 physicians per 1,000 AHCCCS children. For dentists accepting AHCCCS patients, the regional rate was 3 dentists per 1,000 AHCCCS children compared to 16-17 dentists per 1,000 AHCCCS children statewide.

In Table 6, we compared the distance that regional and statewide AHCCCS children needed to travel to the nearest provider type to assist in determining whether the population in the region may have access to care issues based on travel distance. To visit the nearest primary care physician or pharmacy or dentist, 21-29% of AHCCCS children in the region traveled up to one mile for services and another 50-57% traveled one to five miles compared to 56-64% of AHCCCS children statewide who traveled up to one mile and another 26-35% traveled one to five miles. The nearest hospital was one mile away for 10% of regional AHCCCS children, one to five miles away for 66% and more than 10 miles away for 17-18% compared to 11-12% of statewide AHCCCS children who traveled one mile or less, 69% who traveled one to five miles and 10% who traveled 10 miles or more. To visit the nearest behavioral health provider, 38% of regional AHCCCS children traveled one mile and 43% traveled one to five miles; for statewide AHCCCS children, 62-65% traveled one mile to the nearest behavioral health provider and 27-30% traveled one to five miles.

The FOHC sites are Creek Valley Health Clinic in Colorado City. North Countr

<sup>&</sup>lt;sup>19</sup> The FQHC sites are Creek Valley Health Clinic in Colorado City, North Country Healthcare (two in Kingman, one in Bullhead City and one in Lake Havasu City) and River Cities Community Clinic in Bullhead City.

<sup>&</sup>lt;sup>20</sup> The Rural Health Clinics are La Paz Regional Hospital (one in Quartzsite and one in Parker), Parker Medical Center, Parker Walk-in Clinic, San Luis Walk-in Clinic in Parker and Tri-Valley Medical Center in Salome.

Table 4. Supply of Key Health Professionals in Arizona per 1,000 AHCCCS Children, 2017-2019

Provider Type	20:	17	20:	18	2019*	
Provider Type	Number	Rate	Number	Rate	Number	Rate
Total Active Physicians	16,345	70	17,356	74	N/A	N/A
Active Primary Care Physicians <sup>1</sup>	5,396	23	5,598	24	N/A	N/A
Pediatricians <sup>4</sup>	1,214	5	1,257	5	1,293	6
Active Registered and Practical Nurses <sup>2</sup>	N/A	N/A	101,599	433	104,434	445
Dentists <sup>3</sup>	3,796	16	3,903	17	4,012	17

Source: <sup>1</sup> (Association of American Medical Colleges, 2017). (Association of American Medical Colleges, 2019). <sup>2</sup> (National Council of State Boards of Nursing, 2021). <sup>3</sup> (American Dental Association, 2021). <sup>4</sup> (American Board of Pediatrics, 2020); (American Board of Pediatrics, 2019); (American Board of Pediatrics, 2018).

Notes: The rate was calculated using the Arizona population of AHCCCS children birth to age 5 (N = 234,616). National data on pediatricians excluded those who were over age 70 to better control for those who may have been deceased in recent years. <sup>1</sup> Data were from the 2017 and 2019 AMA Physician Masterfiles. Active physicians were federal and non-federal with an Arizona state license who worked at least 20 hours per week. \*Data on active physicians was not available for 2019.

Table 5. Supply of Key Health Professionals in the La Paz/Mohave Region per 1,000 AHCCCS Children, 2017-2019

Provider		2017		2018		19
Provider	Num	Rate	Num	Rate	Num	Rate
Primary Care						
Primary Care – All Licensed Primary Care Physicians <sup>2</sup>		10	82	12	91	14
Physicians accepting AHCCCS <sup>1,2</sup> – Total		8	58	9	63	10
Physicians accepting AHCCCS – Pediatrics		1	8	1	8	1
Physicians accepting AHCCCS – Primary Care		7	50	7	55	8
Physicians with ≥250 AHCCCS patients per year (all ages)		4	24	4	24	4
Behavioral Health – AHCCCS <sup>1</sup>						
Behavioral Health Physician Specialty or Allied Health Professional		8	60	9	63	10
Primary Care with Behavioral Health Services*	7	1	8	1	10	2
Other						
Dentist – accepting AHCCCS <sup>2</sup>		3	21	3	22	3
Hospital <sup>1,3</sup>		1	4	1	4	1
Pharmacy <sup>1,4</sup>	23	3	23	3	23	3

Source: <sup>1</sup>AHCCCS Claims Data, 2021. <sup>2</sup>Arizona Medical Board and Arizona Board of Osteopathic Medical Examiners in Medicine and Surgery, 2021. <sup>3</sup> (Arizona Department of Health Services, 2021). <sup>4</sup>RXOpen dataset, accessed from data.gov, 2020. CHiR was the source for all processing of the AHCCCS and Workforce data.

Notes: The rate was calculated using the regional population of AHCCCS children birth to age 5 in 2019 (N = 6,970 for 2017, N = 6701 for 2018 and N = 6,606 for 2019). Pharmacies that were co-located with hospitals or clinics were not captured in the data. Hospital, and pharmacy historic data was not available, so all numbers are based on most recent data available.

<sup>\*</sup>This includes Federally Qualified Health Clinics and Integrated Clinics. These facilities provide both behavioral health services and physical health services.

Table 6. Percent of AHCCCS Children Grouped by Travel Distance Between Provider and Child's Residence by Provider Type for Region and Arizona, 2017-2019

Dunaidas Tura (86)las	V	0-1 M	liles	1-5 M	1-5 Miles		1iles	10+ M	liles	Unknown**	
Provider Type/Miles	Year	Region	ΑZ	Region	ΑZ	Region	ΑZ	Region	ΑZ	Region	ΑZ
Behavioral Health Specialty or Primary Care with Behavioral Health	2017	38%	62%	43%	30%	4%	3%	10%	2%	4%	4%
	2018	38%	64%	43%	27%	4%	3%	11%	2%	4%	3%
Services*	2019	38%	65%	43%	27%	5%	3%	10%	2%	4%	4%
	2017	24%	62%	54%	29%	8%	3%	13%	4%	1%	2%
Dentist	2018	21%	63%	55%	29%	10%	3%	14%	3%	<1%	3%
	2019	21%	63%	55%	28%	10%	3%	14%	3%	<1%	3%
	2017	10%	11%	66%	69%	7%	9%	17%	10%	<1%	<1%
Hospital	2018	10%	12%	66%	69%	7%	9%	18%	11%	<1%	<1%
	2019	10%	12%	66%	69%	7%	9%	18%	11%	<1%	<1%
	2017	26%	64%	51%	26%	6%	3%	17%	7%	<1%	<1%
Pharmacy	2018	27%	64%	50%	26%	6%	3%	17%	7%	<1%	<1%
	2019	27%	64%	50%	26%	5%	3%	18%	7%	<1%	<1%
	2017	26%	56%	56%	34%	4%	4%	12%	4%	3%	3%
Primary Care Physician	2018	25%	56%	57%	35%	3%	3%	13%	4%	3%	3%
	2019	29%	57%	52%	34%	4%	3%	12%	4%	3%	3%

Source: <sup>1</sup> (AHCCCS Claims Data, 2021). <sup>2</sup> (Arizona Medical Board and Arizona Board of Osteopathic Medical Examiners in Medicine and Surgery, 2021). CHiR was the source for all processing of the AHCCCS and Workforce data.

Notes: See the Approach section for details on this methodology. Pharmacies that were co-located with hospitals or clinics were not captured in the data. Historic data on Hospital and Pharmacy were not available, so all numbers are based on the most recent data available. \*Behavioral Health providers includes primary care providers that offer behavioral health services. \*\*The Unknown column captured children who did not have an exact-match address, so the number of miles to the nearest provider could not be accurately calculated.

#### PRIMARY CARE

Access to primary care is important for the health and well-being of children. Primary care practitioners (PCPs) provide appropriate screenings, treatment and preventive services. When children regularly visit a PCP, they are less likely to visit the emergency department for non-urgent care (Transforming Clinical Practice Initiative, 2019) (Piehl, Clemens, & Joines, 2000).

The inclusion criteria for this indicator were children enrolled in the previous 12 months who had at least one claim with a primary care provider, which includes primary care physicians, nurse practitioners and physician assistants.

Regionally, 81-83% of AHCCCS children ages 25 months to six years had at least one annual PCP visit from 2017 to 2019.

AHCCCS Claims Data, 2021

Statewide, 85-86% of AHCCCS children had at least one PCP visit in Table 7 (Arizona Health Care Cost Containment System, July 2021), and 86-87% of Medicaid children nationally had at least one PCP visit. The regional rates were below the AHCCCS Minimum Performance Standard (MPS)<sup>21</sup> of 84%. The subregions who met or exceeded the MPS for annual PCP visits were Bullhead City (all years), Dolan Springs (2019), Fort Mohave (2019) and Lake Havasu

City (2017 and 2019) (Table 8). Regional AHCCCS children who had annual PCP visits were more likely to be ages 1-2 (88-90%) than ages 3-5 (76-79%) and Hispanic and Latino (84%) than Non-Hispanic or Latino (80-82%) (Figure 6).

Table 7. Arizona and Regional AHCCCS Rates for PCP Visits, 2017-2019


Indicator/Voor	20	17	20	18	2019		
Indicator/Year	Region	Arizona	Region	Arizona	Region	Arizona	
Access to Primary Care	81%	85%	81%	85%	83%	86%	

<sup>&</sup>lt;sup>21</sup> Minimum Performance Standard (MPS) is the minimal expected level of performance by AHCCCS Contractors. AHCCCS-reported rates are the official rates used to determine Contractor compliance with performance requirements. If a Contractor does not achieve the MPS, they will be required to submit a corrective action plan and may be subject to sanctions for each deficient measure.

Table 8. Percent of AHCCCS Children Ages 1 – 5 with a Visit to PCP by Subregion, 2017-2019


Subregion	2017	2018	2019	
Bullhead City	85%	86%	87%	
Colorado City	60%	58%	58%	
Dolan Springs	82%	82%	84%	
Fort Mohave	78%	83%	84%	
Kingman	80%	80%	83%	
Lake Havasu City	85%	82%	85%	
Littlefield	79%	82%	80%	
Parker	68%	71%	55%	
Quartzsite	75%	72%	59%	
Salome	79%	72%	76%	

Figure 6. Percent of AHCCCS Children Ages 1 – 5 with a Visit to PCP by Age Group, Tribal Affiliation, Ethnicity, Race and Year




Percent of children with a visit to primary care practitioner ages 1 - 5


Percent of children with a visit to primary care practitioner ages 1 - 5 by tribal affiliation



Percent of children with a visit to primary care practitioner ages 1 - 5 by ethnicity



Percent of children with a visit to primary care practitioner ages 1 - 5 by race



#### WELL-CHILD VISITS

Well-child visits provide an opportunity for PCPs to examine a child holistically for physical, mental, emotional and social/environmental health. A child's growth and development are tracked during a well-child visit. Screenings, counseling and immunizations take place at well-child visits. PCPs can instill healthy behaviors in children by reinforcing their importance during well-child visits. Parents and caregivers can team up with PCPs to address concerns. Creating a trusted relationship between the PCP and child is important as the child ages and develops, so these visits are beneficial to everyone involved (Moreno, 2018); (Sturgeon, 2015).

This HEDIS indicator assesses whether children who turned 15 months old during the measurement year had one or more well-child visits since birth, categorized by number of visits from one to six or more. A separate HEDIS indicator assesses whether children ages 3-5 had an annual well-child visit.

Regionally in Table 9, 46-53% of AHCCCS children birth to 15 months had six or more well-child visits compared to 53-60% of AHCCCS children statewide and 63-66% of Medicaid children nationally. Approximately 95% of all regional AHCCCS children had at least one well-child visit (Figure 7). The subregions who met or exceeded the AHCCCS statewide rates for six or more well-child visits for AHCCCS children birth to 15 months were Bullhead City (2017 and 2018) and Lake Havasu City (2017 and 2019) (Table 10). Regional AHCCCS children birth to 15 months who had six or more well-child visits were more likely to be Hispanic or Latino (48-61%) than Non-Hispanic or Latino (46-50%) and Asian/Pacific Islander in 2017 and 2019 (58-85%) than other races (Figure 8).

For AHCCCS children ages 3-5 (Table 9), 51-56% of regional children had an annual well-child visit compared to 62-65% of statewide children and 72-74% of Medicaid children ages 3-6 nationally. No subregion met the AHCCCS statewide rates for this indicator (Table 11). Regional AHCCCS children ages 3-5 receiving an annual well-child visit were more likely to be Hispanic or Latino (55-58%) than Non-Hispanic or Latino (49-55%) (Figure 9).

Table 9. Arizona and Regional AHCCCS Rates for Well-Child Visits, 2017-2019

Indicator (Voca	2017		2018		2019	
Indicator/Year	Region	Arizona	Region	Arizona	Region	Arizona
Six or More Well-Child Visits in First 15 Months of Life	46%	53%	49%	58%	53%	60%
Annual Well-Child Visit, Ages 3-5	51%	62%	52%	63%	56%	65%

Number of 95.9% 95.8% 100% Well-Child Visits 93.4% ■ 1 visit Percent of Children 2 visits 80% ■ 3 visits ■ 4 visits 60% 5 visits ■ 6+ visits 40% 20% 0% 2017 2018 2019

Figure 7. Percent of Regional AHCCCS Children by Number of Well-Child Visits Completed During Their First 15 Months by Number of Visits and Year

Table 10. Percent of AHCCCS Children Who Had Six or More Well-Child Visits During Their First 15 Months of Life by Subregion, 2017-2019

Subregion	2017	2018	2019
Bullhead City	54%	59%	58%
Colorado City	DS	16%	DS
Dolan Springs	42%	42% 47%	
Fort Mohave	48%	55%	51%
Kingman	38%	46%	54%
Lake Havasu City	61%	51%	60%
Quartzsite	50%	DS	DS
Salome	DS	DS	55%

Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

Note: Data was suppressed for Littlefield and Parker.

Figure 8. Percent of AHCCCS Children Who Had Six or More Well-Child Visits During Their First 15 Months of Life by Race, Ethnicity, Tribal Affiliation and Year




Table 11. Percent of AHCCCS Children Ages 3-5 with a Well-Child Visit by Subregion, 2017-2019

Subregion	2017	2018	2019
Bullhead City	56%	56%	59%
Colorado City	40%	36%	37%
Dolan Springs	54%	58%	54%
Fort Mohave	47%	50%	58%
Kingman	49%	54%	55%
Lake Havasu City	54%	49%	57%
Littlefield	57%	49%	62%
Parker	33%	42%	DS
Quartzsite	50%	43%	39%
Salome	48%	51%	59%

Percent of children ages 3-5 with well-child visit by race Percent of children ages 3-5 with well-child visit by ethnicity 50.8% 2017 Asian/Pacific Islander 2017 Hispanic or Latino 55.5% 59.0% Black Not Hispanic or Latino 49.4% Caucasian/White 51.6% 2018 Hispanic or Latino 54.5% 30.4% Native American 51.3% Not Hispanic or Latino 51.6% Other/Unknown 58.2% 2019 Hispanic or Latino 2018 Asian/Pacific Islander 57.7% 54.5% Not Hispanic or Latino 52.8% Black 0% 20% 40% 60% 80% 100% Percent of Children 52.3% Caucasian/White Percent of children ages 3-5 with a well-child visit by tribal affiliation 35.2% Native American 2017 2018 2019 Other/Unknown 57.0% 100% 54.5% 2019 Asian/Pacific Islander Percent of Children 80% Black 54.8% 56.1% 60% 51.8% 52.6% Caucasian/White 56.0% 45.7% 43.7% 36.7% 40% 39.2% Native American 57.7% Other/Unknown 20% 0% 20% 40% 60% 80% 100% 0% Percent of Children No Yes No Yes No Yes

Figure 9. Percent of AHCCCS Children Ages 3-5 with a Well-Child Visit by Race, Ethnicity, Tribal Affiliation and Year

# SCREENING FOR LEAD POISONING

Exposure to lead can cause damage to the brain and other vital organs, as well as intellectual and behavioral deficits. Because children who are exposed to lead often have no obvious symptoms, lead poisoning often goes unrecognized. Screening for lead via a capillary or venous lead blood test is an easy way to detect an abnormal blood lead level in children. There is no safe blood lead level. If not found early, exposure to lead and high blood lead levels can lead to irrevocable effects on a child's physical and mental health (Arizona Department of Health Services, 2006); (Arizona Department of Health Services, 2003); (National Center for Environmental Health, 2020).

In Arizona, blood lead results are reportable to the Arizona Department of Health Services (ADHS) for children less than six years old. According to ADHS, children who live in areas designated as high-risk<sup>22</sup> for lead poisoning should receive a blood lead test at 12 and 24 months of age, and older children who have not been previously tested should receive a blood lead test. ADHS reported 61,391 children under age six (14% of children under age 5) were screened in 2019, and 40,773 (66%) of those children lived in high-risk areas. Of the children living in high-risk areas, 29% were screened at 12 months of age, and 19% were screened at 24 months of age. Only 10% of children were screened at both intervals (Arizona Department of Health Services, 2021).

For AHCCCS children being screened for lead poisoning one or more times by their second birthday in, the regional rates increased from 23% in 2017 to 35% in 2019 compared to AHCCCS statewide rates which increased from 32% in 2017 to 35% in 2019 (Table 12). Two subregions met or exceeded the AHCCCS statewide rates for one or more blood lead screenings for lead poisoning by the second birthday: Fort Mohave (2018 and 2019) and Lake Havasu City (2017) (Table 13). Hispanic or Latino AHCCCS children (28-36%) were more likely to be screened for lead poisoning by their second birthday than Non-Hispanic or Latino AHCCCS children (21-35%) (Figure 10).

Table 12. Arizona and Regional AHCCCS Rates for Lead Poisoning Screening, 2017-2019

Indicator/Year	20	17	20	18	20	19
	Region	Arizona	Region	Arizona	Region	Arizona
One or More Tests for Lead Poisoning by Second Birthday	23%	32%	29%	34%	35%	35%

Source: AHCCCS Claims Data, 2021. CHIR is the source for all processing of the AHCCCS data.

<sup>22</sup> Visit <a href="http://www.azhealth.gov/leadmap">http://www.azhealth.gov/leadmap</a> for an interactive map of Arizona neighborhoods to identify those considered to be high-risk.


\_

Table 13. Percent of AHCCCS Children Who Had One or More Tests for Lead Poisoning by Their Second Birthday by Subregion, 2017-2019

Subregion	2017	2017 2018	
Bullhead City	24%	31%	40%
Dolan Springs	23%	32%	42%
Fort Mohave	28%	42%	51%
Kingman	14%	29%	36%
Lake Havasu City	38%	28%	31%

Note: Data was suppressed for Colorado City, Littlefield, Parker, Quartzsite and Salome.

Figure 10. Percent of AHCCCS Children Who Had One or More Tests for Lead Poisoning by Their Second Birthday by Race, Ethnicity, Tribal Affiliation and Year



Note: Data was suppressed for Asian/Pacific Islander in 2017 and 2019, and Native American in 2017.

# WEIGHT ASSESSMENT AND COUNSELING<sup>23</sup>

Childhood obesity has both short-term and long-term effects, so it is important for PCPs to monitor weight problems in children and provide guidance for maintaining a healthy weight and lifestyle. The prevalence of obesity among children aged 2–5 years in 2015-2016 was 14% according to the National Health and Nutrition Examination Survey (Hales, Carroll, Fryar, & Ogden, 2017). For this report, we focused on AHCCCS children ages 3-5.

The regional rates for weight assessment<sup>24</sup> and counseling in Figure 11 showed AHCCCS children in the La Paz/Mohave Region were assessed for weight at 6-7% compared to AHCCCS children statewide who were assessed at rates of 9-19% (Table 14). Rates for nutrition counseling were 1-2% at the regional level versus 4-5% at the state level for AHCCCS children. Physical activity assessments in the region were suppressed, and the state was assessed at rates of <1-1%.<sup>25</sup> The national HEDIS Medicaid rates reported in Table 15 included children ages 3-17, and therefore, were not strictly comparable to the region or state rates for AHCCCS children ages birth to 5.

Table 14. Arizona AHCCCS Rates for Weight Assessment and Counseling, Ages Birth to 5, 2017-2019

Indicator/Year	2017	2017 2018	
BMI Assessment	9%	12%	19%
Nutrition Counseling	4%	5%	5%
Physical Activity Counseling	<1%	1%	1%

<sup>&</sup>lt;sup>23</sup> There was limited reporting in claims data as weight assessment and counseling information was most likely collected in the medical record, so these rates should be interpreted with caution.

<sup>&</sup>lt;sup>24</sup> Under HEDIS, the rates for weight assessment are an evaluation of whether Body mass index (BMI) percentile is accessed and does not determine the absolute BMI value. The diagnosis codes for pediatric BMI included: Z68.51 (< 5th percentile for age), Z68.52 (5th percentile to < 85th percentile for age), Z68.53 (85th percentile to < 95th percentile for age) and Z68.54 (≥ 95th percentile for age).


<sup>&</sup>lt;sup>25</sup> Physical Activity Counseling includes sports physicals which are not provided to children in the early childhood age group.

Table 15. National Medicaid HEDIS Rates for Weight Assessment and Counseling, Ages 3-17 Years, 2017-2019

Indicator/Year	2017	2017 2018	
BMI Assessment	73%	74%	77%
Nutrition Counseling	67%	67%	68%
Physical Activity Counseling	61%	62%	64%

Source: (National Committee for Quality Assurance, 2021).

Figure 11. Percent of AHCCCS Children Ages 3-5 Who Had Weight Assessment and Counseling by Year



Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

Notes: Each of the three items above is a different indicator. Because BMI norms for youth vary with age and sex, this indicator evaluates whether BMI percentile is assessed, rather than an absolute BMI value.

## **DEVELOPMENTAL SCREENING AND DELAY**

During early childhood, children grow and develop at a rapid pace physically and cognitively. Although children develop skills at different times, there are guidelines that define the period when an average child should meet certain developmental milestones. The American Academy of Pediatrics recommends developmental screenings during well-child visits for all children ages 9 months, 18 months, 2 years and 2.5 years (Centers for Disease Control and Prevention). Parents may also notice concerns they have about their child's development and discuss them with their child's health care provider.

Developmental delay occurs when a child does not demonstrate mastery of developmental milestones, and the delay can range from mild to severe. Developmental delays have been found to occur in 10-15% of preschool children (Choo, Agarwal, How, & Yeleswarapu, 2019). The National Health Interview Survey found that from 2015-2018, 18% of U.S. children ages 3-17 years had at least one developmental disability

(Zablotsky & Black, 2020). After being diagnosed with a developmental delay, children should be referred to appropriate behavioral health services.

AHCCCS PCPs use developmental screening tools during 9-month, 18-month and 24-month well-child visits. Developmental screenings are assessed in claims data using billing code CPT 96110. AHCCCS has an active Performance Improvement Project to increase the number of screenings in its eligible populations (Arizona Health Care Cost Containment System, 2021). AHCCCS analyzed its own performance on developmental screenings using several data sources and reported 26% (Median = 42%) of eligible members in acute care screened in 2017 and 30% (Median = 33%) screened in 2018. Rates for AHCCCS children in foster care were 34% and 38% for the same years, respectively. AHCCCS also analyzed the 2018 data for disparities and found disparities in five of Arizona's 15 Counties: Apache, Gila, Navajo, Santa Cruz and Yavapai. Racial disparity was also demonstrated for the American Indian population.

Table 16 showed rates of developmental screenings in AHCCCS children birth to age 5 increased over the report period although still well below the AHCCCS median rates. At the regional level, rates increased 10-16% compared to statewide AHCCCS rates of 10-14%. The subregions where AHCCCS children were most likely to receive developmental screenings were Dolan Springs (2017 and 2019), Kingman (all years) and Lake Havasu City (2017 and 2019) (Table 18). Regional AHCCCS children who received developmental screenings were more likely to be ages 1-2 (28-42%) than age 0 (5-9%) or ages 3-5 (1-4%), and Asian/Pacific Islander in 2017 and 2018 (55-58%) than other races (Figure 12).

Rates of diagnosing developmental delay in AHCCCS children were 2-4% at the regional level compared to 3-5% at the state level for AHCCCS children (Table 16). Regional AHCCCS children who were diagnosed with developmental delay were more likely to be male (3-4%) than female (2-4%), and ages 3-5 (3-5%) than age 0 (1%) or ages 1-2 (2-4%) (Figure 13).

Of those AHCCCS children who were diagnosed with developmental delay, 42-50% of regional AHCCCS children received behavioral health services compared to 47-58% of AHCCCS children statewide (Table 16). The subregions where AHCCCS children who were diagnosed with developmental delay were most likely to receive behavioral health services included Bullhead City (2017 and 2018) and Dolan Springs (2018) (Table 20). Regional AHCCCS children who were diagnosed with developmental delay and received behavioral health services were more likely to be male (50-59%) than female (30-39%), ages 3-5 (45-65%) than ages 1-2 (31-40%) and affiliated with a tribe (60-75%) than not affiliated (41-50%).

<sup>&</sup>lt;sup>26</sup> Due to the limited capture of developmental screenings in claims data alone, these rates should be interpreted with caution.

Table 16. Arizona and Regional AHCCCS Rates for Developmental Screenings and Delay, 2017-2019

Indicator/Year	20	17	2018		2019	
indicatory rear	Region	Arizona	Region	Arizona	Region	Arizona
Developmental Screening, Ages Birth to 5	14%	10%	10%	11%	16%	14%
Diagnosing Developmental Delay, Ages Birth to 5	2%	3%	3%	4%	4%	5%
Developmental Delay and Behavioral Health Services, Ages 3-5	48%	49%	42%	47%	50%	58%

Table 17. Percent of Claims by Provider Type for AHCCCS Children Receiving Developmental Screenings, 2017-2019

Provider Type	20	17	2018		2019	
	Claims Count	Percent of Total	Claims Count	Percent of Total	Claims Count	Percent of Total
Federally Qualified Health Center (FQHC)	<6	DS	13	2%	101	8%
Physician – MD/DO	1,095	84%	443	55%	827	64%
Physician Assistant	55	4%	135	17%	59	5%
Registered Nurse Practitioner	155	12%	208	26%	304	24%

Table 18. Percent of AHCCCS Children Receiving Developmental Screenings, Age 1-2, by Subregion, 2017-2019

Subregion	2017	2018	2019
Bullhead City	13%	14%	25%
Colorado City	DS	DS	14%
Dolan Springs	44%	37%	47%
Fort Mohave	24%	19%	41%
Kingman	46%	42%	50%
Lake Havasu City	48%	34%	58%
Parker	DS	30%	42%
Quartzsite	19%	DS	24%
Salome	DS	DS	38%


Note: Data was suppressed for Littlefield.

Figure 12. Percent of AHCCCS Children Receiving Developmental Screenings by Age Group, Tribal Affiliation, Ethnicity, Race and Year




Percent of children receiving developmental screenings over all


Percent of children receiving developmental screenings over all children ages 1-2 by tribal affiliation



Percent of children receiving developmental screenings over all children ages 1-2 by ethnicity

Percent of children receiving developmental screenings over all children ages 1-2 by race





Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

Notes: Since developmental delay screenings are more likely to take places for those ages 1-2, the other analyses focus on that age group.

Table 19. Percent of AHCCCS Children with a Diagnosed Developmental Delay by Subregion, 2017-2019

Subregion	2017	2018	2019
Bullhead City	2%	2%	1%
Colorado City	DS	2%	3%
Dolan Springs	2%	3%	5%
Fort Mohave	2%	3%	3%
Kingman	3%	5%	7%
Lake Havasu City	3%	3%	4%

Note: Data was suppressed for Littlefield, Parker, Quartzsite and Salome.

Figure 13. Percent of AHCCCS Children with a Diagnosed Developmental Delay by Ethnicity, Sex, Tribal Affiliation, Age Group and Year





Table 20. Percent of AHCCCS Children with a Diagnosed Developmental Delay Who Have Received Behavioral Health Services by Subregion, 2017-2019

Subregion	2017	2018	2019
<b>Bullhead City</b>	76%	61%	56%
Dolan Springs	DS	64%	47%
Fort Mohave	DS	40%	52%
Kingman	41%	38%	50%
Lake Havasu City	49%	43%	54%

Note: Data was suppressed for Colorado City, Littlefield, Parker, Quartzsite and Salome.

Figure 14. Percent of AHCCCS Children with a Diagnosed Developmental Delay Who Have Received Behavioral Health Services by Ethnicity, Race, Sex, Tribal Affiliation, Age Group and Year



Notes: The indicator uses diagnosis code R62.50: Unspecified lack of expected normal physiological development in childhood. Data was suppressed for age 0 in all years and Hispanic or Latino in 2017.

Table 21. Percent of Claims by Provider Type for AHCCCS Children with a Diagnosed Developmental Delay Who Have Received Behavioral Health Services, 2017-2019

	20	17	20	18	2019	
Provider Type	Claims Count	Percent of Total	Claims Count	Percent of Total	Claims Count	Percent of Total
Behavioral Health Outpatient Clinic	902	50%	1,117	46%	1,158	52%
Durable Medical Equipment Supplier	48	3%	100	4%	56	3%
Habilitation Provider	23	1%	77	3%	<6	DS
Hospital	18	1%	18	1%	24	1%
Integrated Clinics	38	2%	196	8%	106	5%
Occupational Therapist	259	14%	379	16%	211	10%
Physical Therapist	26	1%	15	1%	94	4%
Physician – MD/DO	45	3%	36	1%	37	2%
Speech Language Pathology	39	2%	217	9%	170	8%
Speech/Hearing Therapist	368	21%	185	8%	321	14%
Other	26	1%	79	3%	41	2%

#### BEHAVIORAL HEALTH

During the early years of life, the social-emotional development and adaptive functioning of a child changes rapidly and profoundly as their developing brains encounter experiences (National Scientific Council on the Developing Child, 2004). The Adverse Childhood Experiences studies demonstrate how negative early childhood events such as neglect, violence and trauma can lead to behavioral and physical health problems in adulthood like chronic disease, mental illness, and substance abuse (Centers for Disease Control and Prevention, n.d.). However, these effects can be mitigated with proper intervention at the infant and toddler stages by behavioral health services (Arizona Health Care Cost Containment System, 2018). For young children, behavioral health services<sup>27</sup> would likely include day programs, crisis services, rehabilitation services, health promotion, mental health counseling, psychiatric and psychologist services, and various support services.

\_

<sup>&</sup>lt;sup>27</sup> Visit <a href="https://www.azahcccs.gov/Members/AlreadyCovered/coveredservices.html">https://www.azahcccs.gov/Members/AlreadyCovered/coveredservices.html</a> for more details on AHCCCS behavioral health services.

Of AHCCCS children in the region, 12% of children received behavioral services in 2017, 11% received behavioral health services in 2018 and 16% received behavioral health services in 2019.

Pediatric behavioral health providers screen AHCCCS children from birth to age five for emotional, behavioral, and/or developmental needs. A national screening tool assists providers in coordinating services based on the intensity of need and formulating an integrated treatment plan (American Academy of Child and Adolescent Psychiatry, 2006).

In Table 22, 11-16% of regional and statewide AHCCCS children

received behavioral health services. The subregions where AHCCCS children were more likely to receive behavioral health services included Colorado City (2019), Kingman (2019) and Quartzsite (2017) (Table 23). Male AHCCCS children in the region were more likely to receive behavioral health services (13-19%) then females (8-13%) (Figure 15).

Table 22. Arizona and Regional AHCCCS Rates for Behavioral Health Services, Ages 3-5, 2017-2019

Indicator/Voor	2017		2018		2019	
Indicator/Year	Region	Arizona	Region	Arizona	Region	Arizona
Behavioral Health Services, Ages 3-5	12%	11%	11%	12%	16%	16%

Source: AHCCCS Claims Data, 2021. CHIR is the source for all processing of the AHCCCS data.


Table 23. Percent of AHCCCS Children Ages 3-5 Receiving Behavioral Health Services by Subregion, 2017-2019

Subregion	2017	2018	2019
Bullhead City	15%	13%	12%
Colorado City	DS	DS	24%
Dolan Springs	8%	10%	15%
Fort Mohave	9%	8%	10%
Kingman	14%	13%	21%
Lake Havasu City	11%	10%	16%
Quartzsite	19%	13%	12%

Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

Note: Data was suppressed for Littlefield, Parker and Salome.

Figure 15. Percent of AHCCCS Children Ages 3-5 Receiving Behavioral Health Services by Sex, Tribal Affiliation, Ethnicity, Race and Year



Note: Data was suppressed for Asian/Pacific Islander in all years.

Table 24. Percent of Claims by Provider Type for AHCCCS Children Ages 3-5 Receiving Behavioral Health Services, 2017-2019

	2017		2018		2019	
Provider Type	Claims Count	Percent of Total	Claims Count	Percent of Total	Claims Count	Percent of Total
Behavioral Health Outpatient Clinic	7,074	69%	5,495	64%	6,153	62%
Durable Medical Equipment Supplier	66	1%	117	1%	57	1%
Habilitation Provider	473	5%	324	4%	29	<1%
Hospital	77	1%	108	1%	103	1%
Integrated Clinics	553	5%	646	7%	559	6%
Occupational Therapist	555	5%	619	7%	578	6%
Physical Therapist	54	1%	52	1%	148	1%
Physician – MD/DO	189	2%	134	2%	178	2%
Speech Language Pathology	159	2%	435	5%	598	6%
Speech/Hearing Therapist	917	9%	527	6%	1,339	14%
Other	117	1%	191	2%	155	2%

#### VISION

Health conditions such as vision problems are detected through regular visits to PCPs. The American Public Health Association estimates that 20% of preschoolers have eye or vision problems (American Public Health Association, 2019). Vision screenings check the appearance of the eyes and detect potential eye problems. Most vision problems are successfully treated when detected early, but many children do not receive adequate vision screenings. A lack of vision care at younger ages can mean higher rates of undetected vision problems, leading to visual impairments that affect a child's development, performance, and quality of life.

Of all Arizona children 0-5 years old, 35% received a vision screening 2019-2020.

(Child and Adolescent Health Measurement Initiative)

Arizona's Eyes on Learning Vision Coalition recommends a vision screening as early as age one during a well-child visit. Other settings that provide vision screening include pediatrician offices,

educational settings and community settings. Children ages 3-5 should have at least one vision screening by a PCP or trained screener during this timeframe. Annual screenings should be provided to children in kindergarten through fourth grade. A vision screening is not necessary for children with certain developmental delays that cause difficulties with language and speech, motor skills, behavior, memory,

learning, or other neurological functions. Instead, eye doctors recommend that all children with these types of delays receive a comprehensive eye exam. (Eyes on Learning, n.d.)

Vision screenings are typically included in AHCCCS' well-child visits according to their vision periodicity schedule and as medically necessary (Arizona Health Care Cost Containment System, 2021). However, the vision screening is not billed as a separate claim when completed during a well-child visit. Therefore, to capture the population of children who received a vision screening, we assumed that AHCCCS children were screened at their annual well-child visit, or they received a separately billable vision screening. Additional analysis showed that there were very few children who received a vision screening and not a well-child visit annually. Given that the claims data did not specify that a vision screening occurred during the well-child visit, these rates should be interpreted with caution and may be an overestimation of actual vision screenings.

Eye exams are completed by optometrists or ophthalmologists, so we captured those using procedure codes for ophthalmological services. We designated the eye exam as a follow-up eye exam if the visit occurred within six months of a vision screening or well-child visit. If a child was diagnosed with a visually significant eye condition during an eye exam and received treatment or additional visits to an optometrist or ophthalmologist for eyeglasses, surgery or other procedures, the rate of treatment was reported under "visually significant eye conditions who receive treatment". To calculate the rate for visually significant eye conditions who receive treatment, the denominator was all AHCCCS children who received an eye exam and had a diagnosis of strabismus, refraction and accommodation, amblyopia, or other eye disorders; and of those AHCCCS children with an eye condition, the numerator included children who were treated for the eye condition.

In the La Paz/Mohave Region, 45-48% of AHCCCS children received an annual vision screening or well-child visit compared to 43-47% of AHCCCS children statewide (Table 25). Regional AHCCCS children who received an annual vision screening or well-child visit were more likely to be ages 1-2 (69-74%) than ages 3-5 (51-54%) and Hispanic or Latino (49-53%) than Non-Hispanic or Latino (44-47%) (Figure 16). The subregions where AHCCCS children were more likely to receive an annual vision screening or well-child visit included Bullhead City (all years), Dolan Springs (all years), Fort Mohave (all years), Kingman (all years) and Lake Havasu City (2017) (Table 26).

Eye exams were conducted much less frequently, ranging 5-7% annually at the regional level and 4-5% at the state level for AHCCCS children (Table 25). Regional AHCCCS children ages 3-5 (10-11%) were more likely to receive an eye exam than ages 1-2 (2-4%) (Figure 17). Follow-up eye exams were conducted on AHCCCS children in the region and statewide at rates of 4-5%. AHCCCS children with visually significant eye conditions received treatment at rates of 53-67% regionally compared to 54-60% statewide.

Table 25. Arizona and Regional AHCCCS Rates for Vision, 2017-2019

Indicator/Year	2017		2018		2019	
indicator/ rear	Region	Arizona	Region	Arizona	Region	Arizona
Vision Screening or Well-Child Visit	46%	44%	45%	43%	48%	47%
Eye Exams	5%	4%	5%	4%	7%	5%
Follow-up Eye Exams after Vision Screening or Well-Child Visit	5%	4%	5%	5%	5%	4%
Visually Significant Eye Conditions Who Receive Treatment	53%	54%	55%	56%	67%	60%

Table 26. Percent of AHCCCS Children Receiving Vision Screening or Well-Child Visit by Subregion, 2017-2019

Subregion	2017	2018	2019
Bullhead City	51%	52%	55%
Colorado City	35%	33%	27%
Dolan Springs	47%	49%	47%
Fort Mohave	46%	46%	51%
Kingman	46%	48%	49%
Lake Havasu City	47%	39%	44%
Littlefield	23%	25%	46%
Parker	29%	20%	20%
Quartzsite	33%	28%	26%
Salome	19%	18%	38%

Figure 16. Percent of AHCCCS Children Receiving Vision Screening or Well-Child Visit by Age Group, Tribal Affiliation, Ethnicity, Race and Year




Table 27. Percent of AHCCCS Children Receiving Eye Exams by Subregion, 2017-2019

Subregion	2017	2018	2019
Bullhead City	6%	6%	7%
Colorado City	8%	11%	6%
Dolan Springs	3%	2%	6%
Fort Mohave	5%	5%	6%
Kingman	4%	5%	6%
Lake Havasu City	7%	6%	8%

Note: Data was suppressed for Littlefield, Parker, Quartzsite and Salome.




Figure 17. Percent of AHCCCS Children Receiving Eye Exams by Age Group, Tribal Affiliation, Ethnicity, Race and Year

Notes: Eye exams are performed by an optometrist or ophthalmologist. Data was suppressed for Asian/Pacific Islander for all years and Black in 2018.

#### **HEARING**

Most children begin hearing sounds at birth and learn to speak over time by imitating the sounds around them (NIDCD Information Clearinghouse, 2011). The National Institute on Deafness and Other Communication Disorders reports that around two or three out of every 1,000 children are born deaf or hard-of-hearing in the United States, and more lose their hearing later in childhood (NIDCD Information Clearinghouse, 2011). To detect hearing loss early, every state conducts universal newborn hearing screenings before a baby is discharged from a hospital or birthing center. If hearing loss is indicated, parents will be referred to an audiologist to conduct more comprehensive hearing testing and determine appropriate intervention services. For children diagnosed with hearing loss, early intervention helps children develop better language and communication skills.

Arizona strives to screen all infants before 1 month of age. Infants who do not pass the initial hearing screen and a rescreening, should be evaluated further to confirm or diagnose hearing loss before 3 months of age. Infants diagnosed with permanent hearing loss should receive intervention services before 6 months of age (Arizona Department of Health Services, n.d.). This report included available data on hearing screenings along with comprehensive hearing testing, evaluation and assessment which were termed "additional audiology services.

Around 99% (82,035) of all Arizona infants received a newborn hearing screening in 2017 (Arizona Health Care Cost Containment System, 2018) which was slightly higher than the national rate of 98% (National Center on Birth Defects and Developmental Disabilities, 2019). Less than 1% of all Arizona infants were diagnosed with permanent hearing loss, and of those, 42% were diagnosed before three months of age (Arizona Health Care Cost Containment System, 2018). Nationally, 10% of infants were diagnosed with permanent hearing loss, and of those, approximately 74% were diagnosed before three months of age (National Center on Birth Defects and Developmental Disabilities, 2019). Additional audiology services were provided to 4-9% of AHCCCS children under age one in the La Paz/Mohave Region compared to 9-12% of AHCCCS children statewide (Table 28). AHCCCS children under age one had higher rates of additional audiology services in Fort Mohave for all years (Table 29) than other subregions in the La Paz/Mohave Region.

Hearing screenings were provided to 10-11% of AHCCCS children ages 1-5 in the region compared to 20-28% of AHCCCS children statewide. Regional AHCCCS children ages 1-5 were more likely to receive hearing screenings included Dolan Springs (2017) and Lake Havasu City (2018 and 2019) (Table 32). Male regional AHCCCS children ages 1-5 were more likely to receive hearing screenings (12-14%) than females (8-9%) (Figure 19). The provision of additional audiology services ranged 72-88% of AHCCCS children in the region while statewide AHCCCS children's rates decreased from 68% to 57% over the same period. The subregions where AHCCCS children ages 1-5 were more likely to receive additional audiology services included Dolan Springs (2017 and 2018), Fort Mohave (2019), Kingman (2017 and 2019) and Lake Havasu City (all years) (Table 33).

Table 28. Percent of AHCCCS Statewide and Regional Hearing Results, 2017-2019

Indicator / Year	2017		2018		2019	
indicator / Year	Region	Arizona	Region	Arizona	Region	Arizona
Additional Audiology Services Under Age One	9%	11%	9%	12%	4%	9%
Hearing Screening Ages 1-5	11%	20%	11%	22%	10%	28%
Additional Audiology Services for those Screened, Ages 1-5	75%	68%	72%	66%	88%	57%


Table 29. Percent of AHCCCS Children Under Age One Who Had Audiology Services of All Children by Subregion, 2017-2019

Subregion	2017	2018	2019
Bullhead City	15%	14%	5%
Colorado City	9%	DS	DS
Fort Mohave	32%	17%	11%
Kingman	6%	5%	3%
Lake Havasu City	DS	8%	DS

Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

Note: Data was suppressed for Dolan Springs, Littlefield, Parker, Quartzsite and Salome.

Figure 18. Percent of AHCCCS Children Under 1 Year of Age Who Had Audiology Services by Sex, Tribal Affiliation, Ethnicity and Year



Note: Data was suppressed for Hispanic or Latino in 2019, tribal affiliation in 2018 and 2019.

Table 30. Percent of Claims by Provider Type for AHCCCS Children Under Age One Who Had Audiology Services, 2017-2019

	2017		2018		2019	
Provider Type	Claims Count	Percent of Total	Claims Count	Percent of Total	Claims Count	Percent of Total
Audiologist	8	14%	9	20%	<6	DS
Physician – MD/DO	32	55%	25	56%	12	52%
Registered Nurse Practitioner	18	31%	9	20%	<6	DS

Table 31. Percent of Claims by Provider Type for AHCCCS Children Ages 1-5 Receiving Hearing Screening Tests, 2017-2019

	2017		2018		2019	
Provider Type	Claims Count	Percent of Total	Claims Count	Percent of Total	Claims Count	Percent of Total
Audiologist	82	55%	78	52%	30	21%
Physician – MD/DO	66	44%	71	47%	100	70%

Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.


Table 32. Percent of AHCCCS Children Ages 1-5 Receiving Hearing Screening Tests by Subregion, 2017-2019

Subregion	2017	2018	2019
Bullhead City	6%	11%	9%
Colorado City	5%	DS	5%
Dolan Springs	16%	11%	4%
Fort Mohave	11%	12%	9%
Kingman	13%	11%	6%
Lake Havasu City	14%	17%	24%

Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

Note: Data was suppressed for Littlefield, Parker, Quartzsite and Salome.

Figure 19. Percent of AHCCCS Children Ages 1-5 Receiving Hearing Screening Tests by Age Group, Tribal Affiliation, Ethnicity, Race, Sex and Year



Note: Data was suppressed for Asian/Pacific Islander in all years, Black in 2018, Native American in 2017 and 2019, and tribal affiliation in 2017.

Table 33. Percent of AHCCCS Children Ages 1-5 Screened for Hearing and Who Had Additional Audiology Services by Subregion, 2017-2019

Subregion	2017	2018	2019
Bullhead City	65%	60%	65%
Dolan Springs	75%	78%	DS
Fort Mohave	61%	53%	71%
Kingman	70%	58%	90%
Lake Havasu City	90%	100%	100%

Note: Data was suppressed for Colorado City, Littlefield, Parker, Quartzsite and Salome.

Figure 20. Percent of AHCCCS Children Ages 1-5 Screened for Hearing and Who Had Additional Audiology Services by Age Group, Ethnicity, Sex and Year




Table 34. Percent of Audiology Service Claims by Provider Type for AHCCCS Children Ages 1-5 Screened for Hearing and Who Had Additional Audiology Services, 2017-2019

Provider Type	2017		2018		2019	
	Claims Count	Percent of Total	Claims Count	Percent of Total	Claims Count	Percent of Total
Audiologist	62	42%	49	35%	32	22%
Physician – MD/DO	84	56%	84	60%	107	73%
Other	<6	DS	8	6%	8	5%

### **ORAL HEALTH**

Oral health concerns our teeth, gums, and oral-facial system that includes the ability to smile, speak, chew and other senses. Daily brushing and flossing of our teeth and gums demonstrates good oral hygiene, but it is not enough to maintain good oral health. We also need good nutrition, proper management of other health conditions, access to dental care, and extra help when there is a genetic predisposition to oral health conditions or special health care needs.

Oral health is a key indicator of overall health, well-being and quality of life.

Unfortunately, tooth decay has become a chronic disease in children. The CDC reports that 20% of children ages 5-11 have at least one untreated cavity, and children in low-income

families are twice as likely to have cavities than children in higher-income families (Dye, Xianfen, & Beltrán-Aguilar, 2012). Cavities can be prevented by applying a fluoride varnish to primary and permanent teeth every six months, drinking fluoridated tap water, brushing with a fluoride toothpaste, and applying dental sealants. Children should have regular visits to the dentist, beginning before their first birthday, for early identification and management of problems (Enany, n.d.). This report focuses on dental visits for ages 1-5.

In La Paz/Mohave, 51-56% of AHCCCS children had at least one annual dental visit compared to 51-53% of AHCCCS children statewide (Table 35). Neither the region nor the state met the AHCCCS MPS of 60% for annual dental visits for ages 2-20 (Table 36). Colorado City and Kingman subregions met or exceeded the AHCCCS MPS in 2019 (Table 38). Regional AHCCCS children with at least one annual dental visit were more likely to be ages 3-5 (61-65%) than ages 1-2 (36-41%) and Hispanic or Latino (54-58%) than Non-Hispanic or Latino (51-55%) (Figure 21).

Two preventative care dental visits are recommended annually for children. In the region, 22-27% of AHCCCS children received two preventative care dental visits compared to 18-20% of AHCCCS children statewide (Table 35); and 28% of AHCCCS children had one preventative care dental visit (Figure 22). In total, around half of regional AHCCCS children had at least one preventative care dental visit per year. Fluoride varnish was applied to 45-52% of AHCCCS children in the region compared to 47-49% of AHCCCS children statewide (Table 35). The subregions that met or exceeded the AHCCCS statewide rates for

AHCCCS children having received a fluoride varnish application included Bullhead City (all years), Dolan Springs (all years), Fort Mohave (all years) and Kingman (all years) (Table 40). Regional AHCCCS children who had a fluoride varnish application were more likely to be ages 3-5 (55-60%) than ages 1-2 (30-38%), Hispanic or Latino (49-55%) than Non-Hispanic or Latino (44-51%) and Asian/Pacific Islander (50-57%) (Figure 23).

Table 35. Percent of Statewide and Regional AHCCCS Children Oral Health Visits for Ages 1-5, 2017-2019

Type of Visit / Year	20	17	2018		2019	
	Region	Arizona	Region	Arizona	Region	Arizona
Any Annual Dental Visit	51%	51%	52%	52%	56%	53%
Preventative Care Dental Visit Twice Annually	22%	18%	23%	19%	27%	20%
Fluoride Varnish Application	45%	47%	46%	48%	52%	49%

Source: AHCCCS Claims Data, 2021. CHIR is the source for all processing of the AHCCCS data.

Table 36. AHCCCS Statewide Contractor Rate of Performance on Annual Dental Visits for Ages Two to 20 Years, 2017-2019

Contractor	2017	2018	2019	Minimum Performance Standard
AHCCCS Complete Care	61%	61%	60%	60%
Comprehensive Medical and Dental Program	74%	75%	60%	60%
KidsCare	74%	74%	76%	60%

Source: (Health Services Advisory Group, 2021); (Health Services Advisory Group, 2019) (Health Services Advisory Group, 2020).

Table 37. Percent of AHCCCS Claims by Provider Type for Children Ages 1-5 With at Least One Annual Dental Visit, 2017-2019


	2017		2018		2019	
Provider Type	Claims Count	Percent of Total	Claims Count	Percent of Total	Claims Count	Percent of Total
Dentist	5,122	100%	4,972	100%	5,390	100%
Federally Qualified Health Center (FQHC)	22	<1%	11	<1%	12	<1%

Table 38. Percent of AHCCCS Children Ages 1-5 With at Least One Annual Dental Visit by Subregion, 2017-2019

Subregion	2017	2018	2019
Bullhead City	51%	56%	56%
Colorado City	60%	50%	53%
Dolan Springs	58%	50%	58%
Fort Mohave	47%	48%	52%
Kingman	57%	56%	61%
Lake Havasu City	46%	50%	53%
Littlefield	41%	42%	41%
Parker	30%	30%	20%
Quartzsite	39%	32%	30%
Salome	45%	37%	31%

Notes: This indicator includes any claim with an associated dental procedure code (CDT).

Figure 21. Percent of AHCCCS Children Ages 1-5 With at Least One Annual Dental Visit by Age Group, Tribal Affiliation, Ethnicity, Race and Year



Notes: This indicator includes any claim with an associated dental procedure code (CDT).

100%

| 50.7% | 54.6% | 1 visit | 2 or more visits | 1 visit | 1

Figure 22. Percent of AHCCCS Children Ages 1-5 With One and Two Preventative Care Dental Visits in a Year

Notes: This indicator is called a preventative care dental visit and includes the following procedures: D0120 periodic oral evaluation, D0150 comprehensive oral evaluation and D0145 oral evaluation for patient under 3 years of age and counseling with primary caregiver.

Table 39. Percent of Claims by Provider Type for AHCCCS Children Ages 1-5 Receiving Fluoride Varnish, 2017-2019

	2017		2018		2019	
Provider Type	Claims Count	Percent of Total	Claims Count	Percent of Total	Claims Count	Percent of Total
Dentist	3,333	95%	3,268	94%	3,720	94%
Physician – MD/DO	95	3%	111	3%	155	4%
Other	66	2%	95	3%	84	2%

Table 40. Percent of AHCCCS Children Ages 1-5 Receiving Fluoride Varnish by Subregion, 2017-2019

Subregion	2017	2018	2019
Bullhead City	50%	55%	57%
Colorado City	40%	38%	38%
Dolan Springs	54%	49%	54%
Fort Mohave	47%	52%	57%
Kingman	52%	51%	58%
Lake Havasu City	29%	32%	40%
Littlefield	29%	33%	29%
Parker	21%	23%	15%
Quartzsite	30%	24%	27%
Salome	38%	32%	37%

Percent of children receiving flouride varnish ages 1-5 by age group Percent of children receiving flouride varnish ages 1-5 by tribal affiliation 2017 2018 2019 2017 2018 2019 100% 100% Percent of Children Percent of Children 80% 80% 59.9% 54.9% 54.8% 60% 60% 52.2% 46.4% 46.3% 45.0% 41.8% 41.6% 38.4% 40% 40% 33.3% 30.3% 20% 20% 0% 0% 1-2 3-5 1-2 3-5 3-5 1-2 No Yes No Yes No Yes Percent of children receiving flouride varnish ages 1-5 by ethnicity Percent of children receiving flouride varnish ages 1-5 by race 2017 Asian/Pacific Islander 52.1% 2017 Hispanic or Latino 48.5% Black 47.2% Caucasian/White 45.3% Native American 42.6% Not Hispanic or Latino 43.6% Other/Unknown 40.2% 2018 Asian/Pacific Islander 50.0% Hispanic or Latino 50.5% Black 51.4% Caucasian/White 46.7% 47.2% Native American Not Hispanic or Latino 45.0% Other/Unknown 41.8% 2019 Asian/Pacific Islander 56.8% Hispanic or Latino 54.8% 2019 Black 48.1% Caucasian/White 53.8% 40.2% Native American Not Hispanic or Latino 50.6% 41.9% Other/Unknown 0% 20% 40% 60% 80% 100% 0% 20% 40% 60% 80% 100% Percent of Children Percent of Children

Figure 23. Percent of AHCCCS Children Ages 1-5 Receiving Fluoride Varnish by Age Group, Tribal Affiliation, Ethnicity, Race and Year

#### **IMMUNIZATIONS**

Childhood vaccines protect children from many serious and potentially life-threatening diseases such as diphtheria, measles, meningitis, polio, tetanus and whooping cough, at a time in their lives when they are most vulnerable to disease. Approximately 300 children in the United States die each year from vaccine preventable diseases (HHS Office of Disease Prevention and Health Promotion, 2021). Immunizations are essential for disease prevention and are a critical aspect of preventable care for children. Vaccination coverage must be maintained to prevent a resurgence of vaccine-preventable diseases.

The Centers for Medicare and Medicaid Services measures the quality of immunizations through a core indicator of childhood immunization status which is also used by HEDIS. The measure calculates a rate for certain vaccines recommended by a child's second birthday (National Quality Forum, 2017):

- Percent of children who have completed the following schedules: four diphtheria, tetanus and acellular pertussis (DTaP); three polio (IPV); one measles, mumps and rubella (MMR); three haemophilus influenza type B (HiB); three hepatitis B (HepB); one chicken pox (VZV); four pneumococcal conjugate (PCV); one hepatitis A (HepA); two or three rotavirus (RV); two influenza (flu).
- Percent of children who have completed all vaccine courses combined: Combo 10.
- Percent of children who have completed Combo 3: four diphtheria, tetanus and acellular
  pertussis (DTaP); three polio (IPV); one measles, mumps and rubella (MMR); three haemophilus
  influenza type B (HiB); three hepatitis B (HepB); one chicken pox (VZV); four pneumococcal
  conjugate (PCV).

AHCCCS measures childhood immunization completion rates with each of its contractors biennially using the core measure. AHCCCS children's immunization status in Table 42 is the percent of AHCCCS children who have completed each indicated vaccine course by their second birthday, recorded in AHCCCS claims only. These rates were substantially lower than AHCCCS' published statistics in Table 41 due to the limitation of claims data and should be interpreted with caution. AHCCCS declared that claims data does not have the greatest level of detail as claims are not always reported for immunizations, particularly in school settings. To accurately capture immunization rates in AHCCCS' published statistics, AHCCCS uses data from medical records and from the Arizona State Immunization Information System (ASIIS), which is maintained by the Arizona Department of Health Services.

AHCCCS reported that statewide childhood immunization completion rates met or exceeded the national mean rates for three immunizations: DTaP, Hep A and Combo 3 (Arizona Health Care Cost Containment System, 2018) (Table 41). Several barriers to immunizations remained, such as the spread of misinformation about vaccines and parental hesitancy. The rate of exemptions from immunizations increased statewide as nearly 6% of kindergarteners had a Personal Beliefs Exemption in place since the 2017-2018 school year (Arizona Department of Health Services, 2021).

Table 41. AHCCCS Statewide Aggregate Immunization Completion Rates by Two Years Old, FFY 2016

Immunizations	FFY 2016 (period ending 9/30/2017)	HEDIS Medicaid Mean	AHCCCS Minimum Performance Standard
DTaP	79%	77%	85%
Polio	88%	89%	91%
MMR	89%	90%	91%
HiB	87%	88%	90%
Нер В	87%	88%	90%
VZV	88%	89%	88%
PCV	76%	77%	82%
Нер А	88%	84%	40%
RV	61%	69%	60%
Flu	40%	45%	45%
Combo 3	71%	70%	68%

Source: (Arizona Health Care Cost Containment System, 2018).

Note: The rows shaded green are the childhood immunization rates that met or exceeded the national median rates.

Table 42. Percent of Statewide and Regional AHCCCS Children Immunization Status, from AHCCCS Claims Data Only, 2017-2019

	2017		20	2018		19
Immunizations	Region	Arizona	Region	Arizona	Region	Arizona
DTaP	15%	30%	32%	38%	48%	52%
Polio	19%	40%	43%	51%	64%	66%
MMR	47%	57%	71%	72%	71%	76%
HiB	22%	44%	46%	56%	63%	69%
Нер В	8%	13%	13%	18%	24%	21%
VZV	47%	57%	71%	72%	71%	76%
PCV	2%	18%	25%	31%	50%	52%
Нер А	62%	65%	71%	75%	71%	78%
RV	12%	31%	30%	39%	50%	51%
Flu	10%	19%	26%	31%	25%	34%
Combo 3	0%	4%	6%	10%	16%	15%
Combo 10	0%	2%	2%	4%	5%	7%

Notes: Rates only include immunizations recorded in AHCCCS claims, this is likely an undercount of immunization rates. Rows shaded pink are for comparing with Table 41.

#### MATERNAL PRENATAL AND POSTPARTUM CARE

Research has shown that the health of women before pregnancy and after delivery significantly impacts the health of their babies; therefore, it is important to focus on women's preconception health, prenatal care, postpartum care and beyond (Healthy People 2030).

"Women who do not seek prenatal care are three times as likely to deliver a low birth weight infant."

(NICHD - Eunice Kennedy Shriver National Institute of Child Health and Human Development, 2017) For pregnant women, prenatal care is essential for a healthy pregnancy and reducing the complications that can lead to poor birth outcomes for mother and child. Prenatal care involves regular visits to a health care provider to monitor the mother's health and health of the developing fetus, and this care should begin as

early as possible in the pregnancy and continue until delivery. Prenatal care can identify problems or complications and take steps to manage them (NICHD - Eunice Kennedy Shriver National Institute of Child Health and Human Development, 2017). The American Academy of Pediatrics and the American College of Obstetricians and Gynecologists recommend that a woman with an uncomplicated pregnancy be examined at least once in the first trimester for prenatal care. Appropriate perinatal services and education are crucial components of a healthy birth.

The period of up to 60 days following childbirth is called the postpartum period for AHCCCS women. Preexisting health conditions, social determinants, and newly developed conditions contribute to maternal morbidity and mortality during this period. Health care providers consider the postpartum period to be critical to the health and well-being of both mother and child, so postpartum care should not be considered as optional. Yet, research has shown that nearly 40% of women in the United States have gone without a single postpartum visit (American College of Obstetricians and Gynecologists, 2018).

In La Paz/Mohave, 94-95% of pregnant women began prenatal care in the first trimester compared to 84-86% of AHCCCS women statewide (Table 43), which were both above the Healthy People 2030 target rate of 81%<sup>28</sup>. For postpartum care, 96-97% of regional AHCCCS women had at least one postpartum visit compared to 88-89% of AHCCCS women statewide and 64-75% of Medicaid women nationally. Most subregions met or exceeded the Healthy People 2030 target rate for timely prenatal care in all years (Table 44) and postpartum visits (Table 45). The demographic breakdowns for AHCCCS women's prenatal and postpartum care showed high rates, ranging 90-100% (Figure 24 and Figure 25).

<sup>&</sup>lt;sup>28</sup> See <a href="https://health.gov/healthypeople/objectives-and-data/browse-objectives/pregnancy-and-childbirth/increase-proportion-pregnant-women-who-receive-early-and-adequate-prenatal-care-mich-08 for the Healthy People 2030 Prenatal Care Objective.

Table 43. Percent of All AHCCCS Women Who Received Timely Prenatal and Postpartum Care, 2017-2019

Tune of Cave	2017		20	18	2019	
Type of Care	Region	Arizona	Region	Arizona	Region	Arizona
Prenatal Care	95%	84%	95%	86%	94%	85%
Postpartum Care	97%	88%	96%	89%	96%	89%

Table 44. Percent of Deliveries That Received a Prenatal Care Visit While Enrolled in AHCCCS in the First Trimester, on the Enrollment Start Date or Within 42 Days of Enrollment in AHCCCS by Subregion, 2017-2019

Subregion	2017	2018	2019
<b>Bullhead City</b>	93%	96%	96%
Colorado City	95%	91%	88%
Dolan Springs	96%	90%	94%
Fort Mohave	97%	97%	95%
Kingman	97%	94%	94%
Lake Havasu City	98%	96%	96%
Littlefield	93%	91%	100%
Parker	100%	100%	80%
Quartzsite	84%	92%	73%
Salome	75%	100%	58%

Figure 24. Percent of Deliveries That Received a Prenatal Care Visit While Enrolled in AHCCCS in the First Trimester, on the Enrollment Start Date or Within 42 Days of Enrollment in AHCCCS by Age Group, Tribal Affiliation, Ethnicity, Race and Year




Table 45. Percent of AHCCCS Deliveries That Had a Postpartum Visit After Delivery by Subregion, 2017-2019

Subregion	2017	2018	2019
Bullhead City	97%	95%	97%
Colorado City	95%	97%	95%
Dolan Springs	98%	92%	94%
Fort Mohave	98%	98%	96%
Kingman	97%	96%	97%
Lake Havasu City	99%	99%	98%
Littlefield	100%	82%	89%
Parker	100%	100%	90%
Quartzsite	89%	100%	91%
Salome	75%	92%	58%

Percent of deliveries that had a postpartum visit by age group Percent of deliveries that had a postpartum visit by tribal affiliation 2017 2018 2019 2017 2018 2019 98.6% 98.4% 97.3% 97.4% 97.3% 96.8% 97.1% 95.9% 96.1% 95.9% 96.2% 100% 100% 100.0% Percent of Deliveries Percent of Deliveries 80% 80% 60% 60% 40% 40% 20% 20% 0% 0% Adult (20+) Teen (<20) Adult (20+) Teen (<20) Adult (20+) Teen (<20) No Yes No Yes No Yes Percent of deliveries that had a postpartum visit by ethnicity Percent of deliveries that had a postpartum visit by race 2017 Asian/Pacific Islander 94.4% 2017 Hispanic or Latino 97.1% Black 90.5% Caucasian/White 97.5% Native American 100.0% Not Hispanic or Latino 97.3% Other/Unknown 96.7% 2018 Asian/Pacific Islander 100.0% 2018 Hispanic or Latino 93.0% Black 94.4% Caucasian/White 96.0% Native American 100.0% Not Hispanic or Latino 96.9% 95.2% Other/Unknown 2019 Asian/Pacific Islander 100.0% 2019 Hispanic or Latino 94.6% Black 100.0% Caucasian/White 96.6% Native American 95.5% Not Hispanic or Latino 96.6% Other/Unknown 90.1% 0% 20% 60% 80% 100% 0% 20% 60% 80% 100%

Figure 25. Percent of AHCCCS Deliveries That Had a Postpartum Visit After Delivery by Age Group, Race, Ethnicity, Tribal Affiliation and Year

Percent of Deliveries

Percent of Deliveries

#### **HEALTH PLAN PERFORMANCE**

This section provided a selection of health indicators to compare results among the AHCCCS health plans.

Table 47 provided a snapshot of service utilization for regional AHCCCS children enrolled in each AHCCCS health plan. Six regional health plans met or exceeded the AHCCCS statewide aggregate performance and MPS for PCP visits for ages 1-5: Care 1st (2017), Children's Rehabilitative Services (2017 and 2018), Comprehensive Medical and Dental Program (CMDP) (2018 and 2019), DES DD (all years), and Mercy Care Plan (all years) and UnitedHealthcare (2019). Regional performance for at least one well-child visit in the first 15 months was high in at least one year for all reporting health plans. Three health plans met or exceeded the AHCCCS statewide aggregate performance and MPS for annual well-child visits for ages 3-5: CMDP (2018), DES DD (2019) and Mercy Care Plan (2018). CMDP exceeded the MPS for preventative care dental visits for ages 1-5 in 2017 and 2018.<sup>29</sup>

\_

<sup>&</sup>lt;sup>29</sup> The AHCCCS statewide indicator for preventative care dental visits includes ages 2-20 which incorporates a significantly larger number of AHCCCS children than our reporting on ages birth to 5, so the rates should be compared with caution.

Table 46. AHCCCS Statewide Aggregate Health Plan Performance and Minimum Performance Standards, 2017-2019

Year/Indicator	Health Plan Type	One or More Annual PCP Visits, Ages 1-6 (MPS)	Six or More Well-Child Visits in First 15 Months (MPS)	One or More Annual Well- Child Visits, Ages 3-6 (MPS)	One or More Annual Preventative Care Dental Visits, Ages 2- 20 (MPS)
	Acute	83% (84%)	60% (65%)	61% (66%)	61% (60%)
2017	CMDP	92% (84%)	75% (65%)	75% (66%)	74% (60%)
2017	CRS	93% (84%)	49% (65%)	66% (66%)	67% (60%)
	DES DD	89% (84%)	N/A (65%)	53% (66%)	57% (60%)
	Acute	84% (84%)	62% (65%)	61% (66%)	61% (60%)
2018	CMDP	93% (84%)	N/A (65%)	73% (66%)	75% (60%)
2018	CRS	92% (84%)	47% (65%)	64% (66%)	68% (60%)
	DES DD	87% (84%)	N/A (65%)	55% (66%)	57% (60%)
	Acute	84% (87%)	64% (62%)	63% (66%)	60% (60%)
2019	CMDP	92% (87%)	N/A (62%)	75% (72%)	75% (60%)
2019	CRS	N/A (87%)	N/A (62%)	N/A (66%)	N/A (60%)
	DES DD	89% (87%)	N/A (62%)	58% (66%)	53% (60%)

Source: (Health Services Advisory Group, 2019); (Health Services Advisory Group, 2020); (Health Services Advisory Group, 2021); (Health Services Advisory Group, 2019); (Health Services Advisory Group, 2020).

Notes: There was no MPS for DES/DDD for six or more well-child visits in the first 15 months. Cells for which data was not available are indicated by "N/A".

Table 47. Select Regional Indicators by AHCCCS Health Plan, 2017-2019

		One or More PCP		d Visits in Months	One or More Well-Child	One or More Preventative
Health Plan	Year	Visits, Ages 1-5	At Least One Visit*	Six or More Visits	Visits, Ages 3-5	Care Dental Visits, Ages 1-5
	2017	DS	DS	DS	DS	DS
AZ Complete Health	2018	DS	DS	DS	DS	DS
	2019	65%	96%	43%	38%	DS
	2017	83%	DS	DS	64%	42%
Banner University Family Care	2018	79%	78%	DS	53%	51%
Carc	2019	71%	91%	DS	49%	36%
	2017	92%	DS	DS	DS	DS
Care 1st	2018	71%	DS	DS	57%	DS
	2019	82%	98%	53%	55%	42%
Children's Rehabilitative Services (CRS)	2017	89%	91%	DS	63%	48%
	2018	90%	89%	DS	63%	53%
DES Comprehensive	2017	83%	98%	29%	64%	66%
Medical and Dental	2018	90%	100%	25%	69%	70%
(CMDP) Program	2019	91%	98%	51%	55%	59%
DES Developmental	2017	87%	DS	DS	43%	55%
Disability (DD) Long	2018	90%	DS	DS	50%	37%
Term Care (LTC)	2019	91%	DS	DS	90%	59%
	2017	84%	100%	DS	44%	47%
Mercy Care Plan	2018	85%	100%	DS	67%	32%
	2019	94%	DS	DS	DS	30%
	2017	82%	96%	48%	52%	52%
Steward Health Choice AZ	2018	82%	94%	52%	51%	54%
	2019	85%	96%	55%	57%	57%
	2017	81%	98%	48%	49%	49%
United Healthcare	2018	80%	96%	46%	53%	49%
	2019	91%	DS	DS	DS	57%

Notes: The American Indian Health plan was excluded due to low enrollment (1%). Cells shaded green indicate the rate met or exceeded the AHCCCS statewide performance by health plan (Table 48), cells shaded blue indicate the health plan met or exceeded the AHCCCS statewide aggregate performance, cells shaded pink indicate the health plan met or exceeded the MPS, and cells shaded purple indicate the health plan met or exceeded both the MPS and statewide aggregate performance (Table 46).

Table 48. Select AHCCCS Statewide Indicators by Health Plan, 2017-2019

Health Plan	Year	One or More PCP Visits, Ages 1-6	Well-Child Visits in First 15 Months Six or More Visits	One or More Well- Child Visits, Ages 3-6	One or More Preventative Care Dental Visits, Ages 2-20
	2017	N/A	N/A	N/A	N/A
AZ Complete Health	2018	82%	61%	59%	48%
	2019	82%	64%	61%	55%
	2017	N/A	N/A	N/A	N/A
Banner University Family Care	2018	84%	62%	60%	54%
	2019	83%	64%	61%	53%
	2017	83%	66%	64%	62%
Care 1st	2018	86%	67%	67%	65%
	2019	84%	71%	64%	63%
	2017	N/A	N/A	N/A	N/A
Magellan Complete Care	2018	N/A	N/A	N/A	N/A
	2019	67%	N/A	47%	37%
	2017	85%	63%	62%	64%
Mercy Care Plan	2018	86%	66%	63%	64%
	2019	87%	65%	65%	63%
	2017	83%	59%	60%	61%
United Healthcare	2018	84%	61%	61%	62%
	2019	86%	66%	67%	62%

Source: (Health Services Advisory Group, 2019); (Health Services Advisory Group, 2020); (Health Services Advisory Group, 2021); (Health Services Advisory Group, 2019); (Health Services Advisory Group, 2020).

Notes: Cells that did not have available data for that year and/or the health plan was not contracted for that year are indicated by "N/A". Cells shaded blue indicate the rate met or exceeded the AHCCCS statewide aggregate health plan performance rate; cells shaded purple indicate the rate met or exceeded both the AHCCCS statewide aggregate health plan performance rate and the AHCCCS MPS (Table 46). AHCCCS did not measure health plan performance for the indicator of at least one PCP visit by 15 months of age.

#### CONCLUSION

The physical, mental, and emotional health of young children lays the foundation for the rest of their life. La Paz/Mohave had several assets contributing to better health outcomes for young children and women enrolled in AHCCCS from 2017 to 2019, including newborn hearing screenings, immunizations (DTaP, Hepatitis A, Combo 3), vision screenings, and prenatal and postpartum care. These achievements contributed to good health outcomes throughout the region. The areas where needs were identified for AHCCCS women and children included access to care, the supply of health care professionals, PCP visits, developmental screenings, well-child visits, lead poisoning screenings, hearing screenings for ages 1-5 and oral health.

The information in this report can be combined with other available information to create a more comprehensive view of young children and women in the region for regional council planning.

# **REFERENCES**

- American Academy of Child and Adolescent Psychiatry. (2006). *The Early Childhood Service Intensity System (ECSII)*. Retrieved from Arizona Health Care Cost Containment System: https://www.azahcccs.gov/PlansProviders/Downloads/TI/CoreComponents/The%20Early%20Childhood%20Service%20Intensity%20Instrument%20(ESCII)%20.pdf
- American Board of Pediatrics. (2018). *Pediatric Physicians Workforce Data Book, 2017-2018*. Chapel Hill, NC: American Board of Pediatrics. Retrieved from https://www.abp.org/sites/abp/files/pdf/pediatricphysiciansworkforcedatabook2017-2018.pdf
- American Board of Pediatrics. (2019). *Pediatric Physicians Workforce Data Book, 2018-2019.* Chapel Hill, NC: American Board of Pediatrics. Retrieved from https://www.abp.org/sites/abp/files/workforcedata2018-2019.pdf
- American Board of Pediatrics. (2020). *Pediatric Physicians Workforce Data Book, 2019-2020.* Chapel Hill, NC: American Board of Pediatrics. Retrieved from https://www.abp.org/sites/abp/files/pdf/workforcedata2019-2020.pdf
- American College of Obstetricians and Gynecologists. (2018). Optimizing postpartum care. ACOG Committee Opinion No. 736. *Obstet Gynecol, 131*(e140-150). Retrieved from https://www.acog.org/-/media/project/acog/acogorg/clinical/files/committee-opinion/articles/2018/05/optimizing-postpartum-care.pdf
- American Dental Association. (2021). Supply of Dentists in the U.S.: 2001 2020. Chicago: Health Policy Institute, American Dental Association. Retrieved from https://www.ada.org/~/media/ADA/Science%20and%20Research/HPI/Files/HPIData\_SOD\_2020.xlsx?la=en
- American Public Health Association. (2019). Coordinated nationwide approaches to promote eye health and reduce vision impairment. Policy No 20191. Retrieved September 8, 2021, from https://www.apha.org/policies-and-advocacy/public-health-policy-statements/policy-database/2020/01/10/coordinated-nationwide-approaches-to-promote-eye-health-and-reduce-vision-impairment
- Arizona Department of Health Services. (2003, July 8). *Lead A Silent Poison*. Retrieved from Arizona Department of Health Services:

  https://www.azdhs.gov/documents/preparedness/epidemiology-disease-control/lead-poisoning/lead-a-silent-poison-brochure.pdf
- Arizona Department of Health Services. (2006, August 1). What everyone should know about Lead Poisoning. Retrieved from Arizona Department of Health Services:

  https://www.azdhs.gov/documents/preparedness/epidemiology-disease-control/lead-poisoning/lead-poisoning-brochure-english.pdf
- Arizona Department of Health Services. (2021). *Arizona 2019 Annual Report Blood Lead Surveillance Childhood Lead Poisoning Prevention Program.* Phoenix: Office of Environmental Health, Bureau

- of Epidemiology and Disease Control . Retrieved October 11, 2021, from https://azdhs.gov/documents/preparedness/epidemiology-disease-control/lead-poisoning/2019-annual-blood-lead-report.pdf
- Arizona Department of Health Services. (2021). *Arizona Immunization Program*. Retrieved August 30, 2021, from https://www.azdhs.gov/preparedness/epidemiology-disease-control/immunization/index.php#reports-immunization-coverage
- Arizona Department of Health Services. (2021, September). *Consumers Provider & Facility Databases*.

  Retrieved from Medical Facilities Licensing: https://azdhs.gov/licensing/medical-facilities/index.php#consumers-databases
- Arizona Department of Health Services. (n.d.). *Newborn Hearing Screening Early Hearing Detection and Intervention 101*. Retrieved from Arizona Department of Health Services: https://azdhs.gov/preparedness/state-laboratory/newborn-screening/index.php#az-ehdi-home
- Arizona Health Care Cost Containment System. (2018). AHCCCS Complete Care: The Future of Integrated Healthcare Delivery. Retrieved from AHCCCS: https://azahcccs.gov/AHCCCS/Initiatives/AHCCCSCompleteCare/
- Arizona Health Care Cost Containment System. (2018). *Childhood Immunization Completion Rates*.

  Division of Health Care Management, Phoenix. Retrieved from

  https://www.azahcccs.gov/Resources/Downloads/PerformanceMeasures/CYE2017\_ChildhoodI

  mmunizationReport.pdf
- Arizona Health Care Cost Containment System. (2018, August 8). Newborn Screening. Phoenix, AZ.

  Retrieved from

  https://www.azahcccs.gov/Resources/Downloads/Training/Back\_To\_Basics\_Newborn\_Screening.pdf
- Arizona Health Care Cost Containment System. (2018, January 18). Working with birth through five population practice tool. Retrieved August 30, 2021, from https://www.azahcccs.gov/PlansProviders/Downloads/GM/ClinicalGuidanceTools/BirthThrough Five/WorkingWithTheBirthThroughFivePopulationPracticeTool.pdf
- Arizona Health Care Cost Containment System. (2021). Performance improvement project developmental screening interim report. Phoenix. Retrieved from https://www.azahcccs.gov/resources/Downloads/PerformanceImprovementProjects/AHCCCSD evelopmentalScreeningPIPInterimReport.pdf
- Arizona Health Care Cost Containment System. (2021, October). *Policy 430 Attachment A AHCCCS Early and Periodic Screening, Diagnostic and Treatment Periodicity Schedule.* Retrieved from AHCCCS Medical Policy Manual: https://www.google.com/url?client=internal-element-cse&cx=016416319741515762556:nfpz2wqmsus&q=https://www.azahcccs.gov/shared/Downloads/MedicalPolicyManual/400/430\_AttachmentA.docx&sa=U&ved=2ahUKEwjs5f\_b8rL3AhWsJUQIHf4wCCIQFnoECAMQAQ&usg=AOvVaw2PugsSuiAzf

- Arizona Health Care Cost Containment System. (July 2021). AHCCCS Quality Strategy Evaluation CYE 2018 CYE 2020. Phoenix. Retrieved from https://azahcccs.gov/PlansProviders/Downloads/QualityStrategyEvaluationCYE2018-CYE2020.pdf
- Association of American Medical Colleges. (2017). 2017 State Physician Workforce Data Report. Washington, DC: AAMC.
- Association of American Medical Colleges. (2019). 2019 State Physician Workforce Data Report. Washington, DC: AAMC.
- Centers for Disease Control and Prevention. (n.d.). *Adverse Childhood Experiences*. (N. C. Control, Producer) Retrieved from Violence Prevention:

  https://www.cdc.gov/violenceprevention/aces/index.html
- Centers for Disease Control and Prevention. (n.d.). Learn more about your child's development: developmental monitoring and screening. Retrieved August 27, 2021, from https://www.cdc.gov/ncbddd/actearly/pdf/Dev-Mon-and-Screen-English-and-Spanish-P.pdf
- Choo, Y. Y., Agarwal, P., How, C. H., & Yeleswarapu, S. P. (2019). Developmental delay: identification and management at primary care level. *Singapore Medical Journal, 60*(3), 119-123. doi:10.11622/smedj.2019025
- Dye, B. A., Xianfen, L., & Beltrán-Aguilar, E. D. (2012). *Selected oral health indicators in the United States* 2005–2008. Hyattsville, MD: National Center for Health Statistics, Centers for Disease Control and Prevention.
- Enany, E. (n.d.). Small smiles: what you need to know about baby teeth. (A. D. Association, Producer)

  Retrieved August 28, 2021, from mouthhealthy.org: https://www.mouthhealthy.org/en/babies-and-kids/childrens-dental-health
- Eyes on Learning. (n.d.). Why Screening. Retrieved from Eyes on Learning: http://eyesonlearning.org/why-childrens-vision-matters/why-screening/
- Hales, C. M., Carroll, M. D., Fryar, C. D., & Ogden, C. L. (2017). *Prevalence of obesity among adults and youth: United States, 2015–2016.* Hyattsville, MD:: National Center for Health Statistics.
- Health Services Advisory Group. (2019). CYE 2018 Annual Report for Acute Care and CMDP. Phoenix:

  Arizona Health Care Cost Containment System. Retrieved from

  https://www.azahcccs.gov/Resources/Downloads/EQR/2018/CYE\_2018\_External\_Quality\_Revie
  w\_Annual\_Report\_Acute\_Care\_And\_CMDP.pdf
- Health Services Advisory Group. (2019). External Quality Review Annual Report for Arizona Long Term

  Care System Contractors. Phoenix: Arizona Health Care Cost Containment System. Retrieved

  from

  https://azahcccs.gov/Resources/Downloads/EQR/2018/CYE\_2018\_External\_Quality\_Review\_An
  nual Report ALTCS.pdf

- Health Services Advisory Group. (2020). *CYE 2019 Annual Report for 2018 Acute Care and CMDP, RBHAs, and CRS.* Phoenix: Arizona Health Care Cost Containment System. Retrieved from https://www.azahcccs.gov/Resources/Downloads/EQR/2019/CYE2019ExternalQualityReviewAn nualReport-AcuteCMDPRBHACRS.pdf
- Health Services Advisory Group. (2020). External Quality Review Annual Report for Arizona Long Term

  Care System Contractors. Phoenix: Arizona Health Care Cost Containment System. Retrieved from

  https://azahcccs.gov/Resources/Downloads/EQR/2019/CYE2019ExternalQualityReviewAnnualReportALTCS.pdf
- Health Services Advisory Group. (2021). *CYE 2019 Annual Report for AHCCCS Complete Care and CMDP.*Phoenix: Arizona Health Care Cost Containment System. Retrieved October 18, 2021, from https://www.azahcccs.gov/Resources/Downloads/EQR/2020/CYE2020ExternalQualityReviewAnnualReportACCandCMDP.pdf
- Health Services Advisory Group. (2021). External Quality Review Annual Report for Arizona Long Term Care System. Phoenix: Arizona Health Care Cost Containment System. Retrieved from https://azahcccs.gov/Resources/Downloads/EQR/2020/CYE2020ExternalQualityReviewAnnualReportEPDandDDD.pdf
- Healthy People 2030. (n.d.). Pregnancy and Childbirth. Office of Disease Prevention and Health Promotion, Office of the Assistant Secretary for Health, Office of the Secretary, U.S. Department of Health and Human Services. Retrieved August 25, 2021, from https://health.gov/healthypeople/objectives-and-data/browse-objectives/pregnancy-and-childbirth
- HHS Office of Disease Prevention and Health Promotion. (2021, October). *Immunization and Infectious Diseases*. Retrieved from HealthyPeople.gov: https://www.healthypeople.gov/node/3527/data-details.%C2%A0Accessed
- Moreno, M. A. (2018). The Well-Child Visit. *JAMA Pediatr*, *172*(1), 104. doi:10.1001/jamapediatrics.2017.4041
- National Center for Environmental Health. (2020, January). *Health Effects of Lead Exposure*. Retrieved from Centers for Disease Control and Prevention: https://www.cdc.gov/nceh/lead/prevention/health-effects.htm
- National Center on Birth Defects and Developmental Disabilities. (2019, August). Summary of 2017

  National CDC EHDI Data. Retrieved from Centers for Disease Control and Prevention:

  https://www.cdc.gov/ncbddd/hearingloss/2017-data/documents/01-2017-HSFS-Data-Summary.pdf
- National Committee for Quality Assurance. (2021). Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents. Retrieved from HEDIS:

  https://www.ncqa.org/hedis/measures/weight-assessment-and-counseling-for-nutrition-and-physical-activity-for-children-adolescents/

- National Council of State Boards of Nursing. (2021). *Number of Nurses in U.S. and by Jurisdiction*. Retrieved from NCSBN: https://www.ncsbn.org/14283.htm
- National Quality Forum. (2017, January 23). *Childhood Immunization Status*. Retrieved August 30, 2021, from https://www.qualityforum.org/QPS/QPSTool.aspx?m=240#qpsPageState=%7B%22TabType%22%3A1,%22TabContentType%22%3A2,%22ItemsToCompare%22%3A%5B%5D,%22StandardID%22%3A240,%22EntityTypeID%22%3A1%7D
- National Scientific Council on the Developing Child. (2004). *Children's Emotional Development Is Built into the Architecture of Their Brains*. Center on the Developing Child at Harvard University. Retrieved from http://www.developingchild.net
- NICHD Eunice Kennedy Shriver National Institute of Child Health and Human Development. (2017, January 31). *Pre-Pregnancy Care and Prenatal Care*. Retrieved August 27, 2021, from https://www.nichd.nih.gov/health/topics/factsheets/preconceptioncare
- NIDCD Information Clearinghouse. (2011, May). NIDCD Fact Sheet: It's Important to Have Your Baby's Hearing Screened. Retrieved from National Institute on Deafness and Other Communication: https://www.nidcd.nih.gov/sites/default/files/Documents/health/hearing/ItsImportantToHaveY ourBabysHearingScreenedWebFS.pdf
- Piehl, M. D., Clemens, C. J., & Joines, J. D. (2000, Aug). "Narrowing the Gap": decreasing emergency department use by children enrolled in the Medicaid program by improving access to primary care. *Arch Pediatr Adolesc Med*, *154*(8), 791-5. doi:10.1001/archpedi.154.8.791
- Sturgeon, M. (2015, December 15). Parent Plus: Importance of routine pediatrician visits. *AAP News*. The American Academy of Pediatrics. Retrieved from https://www.aappublications.org/news/aapnewsmag/2015/12/15/WellChild121515.full.pdf
- Transforming Clinical Practice Initiative. (2019, December 6). *Reducing Unnecessary Emergency Department Visits*. Retrieved from Centers for Medicare and Medicaid Services: https://innovation.cms.gov/files/x/tcpi-changepkgmod-edvisits.pdf
- Zablotsky, B., & Black, L. (2020). *Prevalence of children aged 3–17 years with developmental disabilities, by urbanicity: United States, 2015–2018.* Hyattsville, MD: National Health Statistics Reports No 139. Retrieved from https://www.cdc.gov/nchs/data/nhsr/nhsr139-508.pdf

## **APPENDIX: DATA SOURCES**

The source of data in all tables, graphs, reports, presentations, and other publications is the Arizona Health Care Cost Containment System (AHCCCS) (2021) unless otherwise noted. CHiR is the source of the calculations, analysis and/or processing of the data.

**AHCCCS** health claims and encounters data. AHCCCS is the state Medicaid provider. The data include health care transactions (paid claims) on all members, patients receiving inpatient, emergency department or other outpatient care in the state. The data layout is already at the individual patient level when received and requires no further manipulation to standardize variables or match patients.

### AHCCCS System

In October 2018, AHCCCS enacted major changes to its care delivery system to integrate physical and behavioral health care under designated health plans for its eligible populations, called AHCCCS Complete Care. Integrated care would result in better coordination among providers in the same network and better health outcomes for AHCCCS enrollees. Under AHCCCS Complete Care, the choice of health plans varies by geographic area, but affected members have the same array of covered services and access to a network of providers (Arizona Health Care Cost Containment System, 2018).

Table 49. AHCCCS Complete Care Health Plans by Geographic Service Area

Geographic Service Area	AHCCCS Complete Care Health Plans
North (Apache, Coconino, Mohave, Navajo and Yavapai Counties)	Care 1st and Health Choice Arizona
Central (Maricopa, Gila and Pinal Counties)	Banner University Family Care, Care 1st, Health Choice Arizona, Arizona Complete Health, Magellan Complete Care, Mercy Care, UnitedHealthcare Community Plan
South (Cochise, Graham, Greenlee, La Paz, Pima, Santa Cruz and Yuma Counties)*	Banner University Family Care, Arizona Complete Health, UnitedHealthcare Community Plan (Pima County only)

<sup>\*</sup>Zip codes 85542, 85192, 85550 are in the South geographic service area.

Other health plans serve specialty populations. AHCCCS members with developmental disabilities who are enrolled in the Department of Economic Security/Division of Developmental Disabilities (DES/DDD) with a Children's Rehabilitative Services designation receive integrated care through their assigned DDD health plan. Arizona Long Term Care members receive services through program contractors.

American Indian members have the choice of enrolling in an AHCCCS Complete Care managed care plan or the American Indian Health Program (AIHP fee-for-service) for integrated care or switch enrollment between the two at any time. AIHP members can also choose care coordination through a Tribal Regional Behavioral Health Authority when available (secondary health plan). American Indian members can receive services at any time from an Indian Health Service facility, or a tribally owned or operated organization (i.e., Tribal 638 providers or Urban Indian Health providers).

American Indian members determined to have a Serious Mental Illness receive behavioral health services from a Regional Behavioral Health Authority but have the option to choose the American Indian Health Program or AHCCCS Complete Care for physical health services.

**Arizona Health Care Workforce- Physicians.** For the provider indicators, we capture the supply of Arizona physicians by specialty using the Arizona Health Care Workforce data set. This data set includes administrative data collected from the Arizona Medical Board and the Arizona Board of Osteopathic Examiners in Medicine and Surgery, the licensing agencies for physicians.