FIRST THINGS FIRST

Navajo/Apache Region



2022 Needs and Assets Supplemental Report:

Children's Access to and Use of Public Health Services

Report Prepared by:

Center for Health Information & Research

Arizona State University

502 E. Monroe St, Suite C320

Phoenix, AZ 85004

(602) 496-2009 | chir@asu.edu / chs.asu.edu/chir

Project Team

- Molly Loughran, MS, Data Science Specialist
- Varnika Angampally, MS, Statistical Programmer
- Nishanth Prathap, MS, Data Science Specialist
- Tameka Sama, MBA, CRA, Center Administrator
- Sruthi Kommareddy, Database Analyst
- Gevork Harootunian, MS, Data Science Consultant
- Meghan Morris, MD, Clinical Consultant
- Anita Murcko, MD, Clinical Consultant

Acknowledgements

The authors gratefully acknowledge the cooperation of First Things First representatives Melissa Affronti, Evaluation Project Director, and Roopa Iyer, Senior Director for Research and Evaluation for their valuable technical support and for the feedback provided by the Regional Needs and Assets (RNA) Health Base Expansion Workgroup along with the RNA Workgroup. Special thanks to Candida Hunter and Kalvina Belin for their skilled administration and management of tribal relations and to Lisa Colling for her assistance with Geocoding and mapping of FTF regions and subregions.



Center for Health Information and Research

TABLE OF CONTENTS

List of Tables	
List of Figures	5
Executive Summary	8
Introduction	16
The Importance of Early Childhood Health	16
Definitions	16
Approach	19
Reporting	20
Data Limitations	21
Navajo/Apahce Regional Results	22
Population and Demographics of Children Enrolled in AHCCCS	23
Health Care Workforce	28
Primary Care	32
Well-Child Visits	35
Screening for Lead Poisoning	41
Weight Assessment and Counseling	44
Developmental Screening and Delay	45
Behavioral Health	53
Vision	57
Hearing	63
Oral Health	69
Immunizations	76
Maternal Prenatal and Postpartum Care	79
Health Plan Performance	85
Conclusion	89
References	90
Amman divu Data Cauraaa	0.5

LIST OF TABLES

Table 1. Number of AHCCCS Children Birth to Age 5 by Year and Sex	23
Table 2. Number of AHCCCS Children Birth to Age 5 by Year and Age Group	23
Table 3. Number of AHCCCS Claims by Provider Type (Billing Entity), 2017-2019	27
Table 4. Supply of Key Health Professionals in Arizona per 1,000 AHCCCS Children, 2017-2019	29
Table 5. Supply of Key Health Professionals in Navajo/Apache Region per 1,000 AHCCCS Children,	2017-
2019	30
Table 6. Percent of AHCCCS Children Grouped by Travel Distance Between Provider and Child's Resident	dence
by Provider Type for Region and Arizona, 2017-2019	31
Table 7. Arizona and Regional AHCCCS Rates for PCP Visits, 2017-2019	32
Table 8. Arizona and Regional AHCCCS Rates for Well-Child Visits, 2017-2019	35
Table 9. Arizona and Regional AHCCCS Rates for Lead Poisoning Screening, 2017-2019	41
Table 10. Arizona AHCCCS Rates for Weight Assessment and Counseling, Ages Birth to 5, 2017-2019)44
Table 11.National Medicaid HEDIS Rates for Weight Assessment and Counseling, Ages 3-17 Years,	2017-
2019	45
Table 12. Arizona and Regional AHCCCS Rates for Developmental Screenings and Delay, 2017-2019	46
Table 13. Percent of Claims by Provider Type for AHCCCS Children Receiving Developmental Scree	nings,
2017-2019	47
Table 14. Percent of Claims by Provider Type for AHCCCS Children with a Diagnosed Developmental	Delay
Who Have Received Behavioral Health Services, 2017-2019	53
Table 15. Arizona and Regional AHCCCS Rates for Behavioral Health Services, Ages 3-5, 2017-2019.	54
Table 16. Percent of Claims by Provider Type for AHCCCS Children Ages 3-5 Receiving Behavioral H	
Services, 2017-2019	
Table 17. Arizona and Regional AHCCCS Rates for Vision, 2017-2019	
Table 18. Percent of AHCCCS Statewide and Regional Hearing Results, 2017-2019	64
Table 19. Percent of Claims by Provider Type for AHCCCS Children Ages 1-5 Receiving Hearing Scre	ening
Tests, 2017-2019	
Table 20. Percent of Audiology Service Claims by Provider Type for AHCCCS Children Ages 1-5 Scr	
for Hearing and Who Had Additional Audiology Services, 2017-2019	
Table 21. Percent of Statewide and Regional AHCCCS Children Oral Health Visits for Ages 1-5, 2017	'-2019
	_
Table 22. AHCCCS Contractor Rate of Performance on Annual Dental Visits for Ages Two to 20 Years,	2017-
2019	
Table 23. Percent of AHCCCS Claims by Provider Type for Children Ages 1-5 With at Least One A	
Dental Visit, 2017-2019	
Table 24. Percent of Claims by Provider Type for AHCCCS Children Ages 1-5 Receiving Fluoride Va	-
2017-2019	
Table 25. AHCCCS Statewide Aggregate Immunization Completion Rates by Two Years Old, FFY 2010	
Table 26. Percent of Statewide and Regional AHCCCS Children Immunization Status, from AHCCCS C	
Data Only 2017-2019	78

Table 27. Percent of All AHCCCS Women Who Received Timely Prenatal and Postpartum Care, 2017-2019
Table 28. AHCCCS Statewide Aggregate Health Plan Performance and Minimum Performance Standards, 2017-2019
Table 29. Select Regional Indicators by AHCCCS Health Plan, 2017-201987
Table 30. Select AHCCCS Statewide Indicators by Health Plan, 2017-201988
Table 31. AHCCCS Complete Care Health Plans by Geographic Service Area95
LIST OF FIGURES
Figure 1. Number of AHCCCS Children Birth to Age 5 by Subregion by Year24
Figure 2. Number of AHCCCS Children Birth to Age 5 by Year and Race24
Figure 3. Number of AHCCCS Children Birth to Age 5 by Year and Ethnicity25
Figure 4. Number of AHCCCS Children Birth to Age 5 by Year and Tribal Affiliation25
Figure 5. Number of AHCCCS Children by Year and Health Plan
Figure 6. Percent of AHCCCS Children Ages 1 – 5 with a Visit to PCP by Subregion by Year33
Figure 7. Percent of AHCCCS Children Ages $1-5$ with a Visit to PCP by Age Group, Tribal Affiliation,
Ethnicity, Race and Year34
Figure 8. Percent of Regional AHCCCS Children by Number of Well-Child Visits Completed During Their
First 15 Months by Number of Visits and Year
Figure 9. Percent of AHCCCS Children Who Had Six or More Well-Child Visits During Their First 15 Months
of Life by Subregion and Year37
Figure 10. Percent of AHCCCS Children Who Had Six or More Well-Child Visits During Their First 15 Months
of Life by Race, Ethnicity, Tribal Affiliation and Year
Figure 11. Percent of AHCCCS Children Ages 3-5 with a Well-Child Visit by Subregion by Year39
Figure 12. Percent of AHCCCS Children Ages 3-5 with a Well-Child Visit by Race, Ethnicity, Tribal Affiliation
and Year40
Figure 13. Percent of AHCCCS Children Who Had One or More Tests for Lead Poisoning by Their Second
Birthday by Subregion by Year42
Figure 14. Percent of AHCCCS Children Who Had One or More Tests for Lead Poisoning by Their Second
Birthday by Race, Ethnicity, Tribal Affiliation and Year
Figure 15. Percent of AHCCCS Children Ages 3-5 Who Had Weight Assessment and Counseling by Year 45
Figure 16. Percent of AHCCCS Children Receiving Developmental Screenings, Age 1-2, by Subregion by
Year47
Figure 17. Percent of AHCCCS Children Receiving Developmental Screenings by Age Group, Tribal
Affiliation, Ethnicity, Race and Year
Figure 18. Percent of AHCCCS Children with a Diagnosed Developmental Delay by Subregion by Year 49
Figure 19. Percent of AHCCCS Children with a Diagnosed Developmental Delay by Ethnicity, Race, Sex,
Tribal Affiliation, Age Group and Year50
Figure 20. Percent of AHCCCS Children with a Diagnosed Developmental Delay Who Have Received
Behavioral Health Services by Subregion by Year51

Figure 21. Percent of AHCCCS Children with a Diagnosed Developmental Delay Who Have Received Behavioral Health Services by Sex and Year
Figure 22. Percent of AHCCCS Children with a Diagnosed Developmental Delay Who Have Received Behavioral Health Services by Age Group and Year
Figure 23. Percent of AHCCCS Children Ages 3-5 Receiving Behavioral Health Services by Subregion by Year54
Figure 24. Percent of AHCCCS Children Ages 3-5 Receiving Behavioral Health Services by Sex, Tribal
Affiliation, Ethnicity, Race and Year55
Figure 25. Percent of AHCCCS Children Receiving Vision Screening or Well-Child Visit by Subregion by Year
Figure 26. Percent of AHCCCS Children Receiving Vision Screening or Well-Child Visit by Age Group, Tribal Affiliation, Ethnicity, Race and Year
Figure 27. Percent of AHCCCS Children Receiving Vision Screening or Well-Child Visit by Subregion by Year
Figure 28. Percent of AHCCCS Children Receiving Eye Exams by Age Group, Tribal Affiliation, Ethnicity, Race and Year
Figure 29. Percent of AHCCCS Children Under 1 Year of Age Who Had Audiology Services by Sex and Year
Figure 30. Percent of AHCCCS Children Ages 1-5 Receiving Hearing Screening Tests by Subregion by Year 65
Figure 31. Percent of AHCCCS Children Ages 1-5 Receiving Hearing Screening Tests by Age Group, Tribal Affiliation, Ethnicity, Race and Year
Figure 32. Percent of AHCCCS Children Ages 1-5 Screened for Hearing and Who Had Additional Audiology Services by Subregion by Year
Figure 33. Percent of AHCCCS Children Ages 1-5 Screened for Hearing and Who Had Additional Audiology Services by Age Group, Tribal Affiliation, Ethnicity, Race and Year
Figure 34. Percent of AHCCCS Children Ages 1-5 With at Least One Annual Dental Visit by Subregion by Year
Figure 35. Percent of AHCCCS Children Ages 1-5 With at Least One Annual Dental Visit by Age Group, Tribal Affiliation, Ethnicity, Race and Year
Figure 36. Percent of AHCCCS Children Ages 1-5 With One and Two Preventative Care Dental Visits in a Year
Figure 37. Percent of AHCCCS Children Ages 1-5 Receiving Fluoride Varnish by Subregion by Year
Figure 39. Percent of Deliveries That Received a Prenatal Care Visit While Enrolled in AHCCCS in the First
Trimester, on the Enrollment Start Date or Within 42 Days of Enrollment in AHCCCS by Subregion by Year
Figure 40. Percent of Deliveries That Received a Prenatal Care Visit While Enrolled in AHCCCS in the First Trimester, on the Enrollment Start Date or Within 42 Days of Enrollment in AHCCCS by Age Group, Tribal Affiliation, Ethnicity, Race and Year

Figure 41. Percent of AHCCCS Deliveries That Had a Postpartum Visit After Delivery by	Subregion by Year
	83
Figure 42. Percent of AHCCCS Deliveries That Had a Postpartum Visit After Delivery by	y Age Group, Race
Ethnicity. Tribal Affiliation and Year	84

EXECUTIVE SUMMARY

Under the direction of First Things First (FTF), the Arizona State University Center for Health Information & Research (CHiR) conducted a regional network analysis of children from birth to age 5 to determine the health assets and health needs in the Navajo/Apache Region. Navajo/Apache consisted of the following subregions: Blue Ridge, Concho, Holbrook, Show Low, Snowflake and Vernon. The main data source was claims data from the Arizona Health Care Cost Containment System (AHCCCS), Arizona's Medicaid agency; therefore, the results presented in this report were for children and mothers who were enrolled in AHCCCS from 2017 to 2019. This population was denoted as AHCCCS children or AHCCCS women.

CHiR and representatives from the FTF Regions, Programs, and Evaluation teams determined priority indicators for this report. AHCCCS children's health was measured in the following categories: primary care and well-child visits, health care workforce, screening for lead poisoning, weight assessment and counseling, developmental health, behavioral health, vision, hearing, oral health, immunizations, maternal prenatal and postpartum care, and health plan performance. Many of the reported indicators were from the Healthcare Effectiveness Data and Information Set (HEDIS)². HEDIS is a performance improvement tool whereby health plans, health care organizations and government agencies submit data on specific health measures. HEDIS uses the collected data to calculate national performance statistics and benchmarks and set standards for measures. HEDIS specifications were applied to the AHCCCS population for each region. Non-HEDIS indicators, which do not have associated benchmarks, were compared to state and national data when possible. The results were displayed by gender, age, race, ethnicity, tribal affiliation³, provider type, and health plan when the data was available and within data suppression guidelines. The results of the analyses are summarized below. When possible, the results are grouped by 1) indicators that met or were above the state average or national HEDIS standards and 2) indicators that did not meet or were below the state average or national HEDIS standards. Other notable findings are also presented that do not have comparison data.

Population and Demographics of Children Enrolled in AHCCCS

There were 234,616 children from birth to age 5 enrolled in AHCCCS statewide from 2017 to 2019. In Navajo/Apache there were 3,077 children enrolled in AHCCCS in 2017, 2,968 children enrolled in 2018 and 2,920 children enrolled in 2019. Of these, male AHCCCS children outnumbered females by 2-4%. There were 6% more infants and toddlers than preschoolers in 2017, with the difference dropping to 1% more in 2019.

¹ Data used in this report covers all AHCCCS members in Arizona, including members living in FTF tribal regions and subregions. Reports for tribal regions and subregions were carried out with specific approval from each tribe. For those tribal regions and subregions who did not give approval, data is included only in aggregate totals for Arizona, and—in the case of a tribal subregion—aggregate totals for the region.

² See https://www.ncqa.org/hedis/

³ Tribal affiliation refers to whether an individual is a member of a federally recognized Arizona tribe and is displayed as a flag (Yes/No) in this report. This information is captured during enrollment in AHCCCS.

Approximately 65% of the AHCCCS children in Navajo/Apache Region lived in three subregions: Holbrook, Show Low and Snowflake. By race, 62-66% of AHCCCS children were reported as Caucasian/White, 20% were Native American, 2-3% were Black and 1% were Asian/Pacific Islander. By ethnicity, 82-83% were reported as Non-Hispanic or Latino and 17-18% were Hispanic or Latino (Figure 3). Affiliation with a tribal community was reported by 23-24% of AHCCCS children annually. Most AHCCCS children in the region were enrolled in three health plans: Steward Health Choice AZ (48-49%), UnitedHealthcare (21-31%) and the American Indian Health Plan (10%). By health claim, 29-32% were submitted by physicians, 13-15% were submitted by pharmacies and 12-13% were submitted by hospitals

Health Care Workforce

The supply of physicians in the United States is tracked by the Association of American Medical Colleges biennially. Arizona had 160 hospitals individually licensed by the state which were subtyped as children, critical access, long term, short term, psychiatric, rehabilitation, transplant and non-participating. The Navajo/Apache Region had one short-term hospital, Summit Healthcare Regional Medical Center in Show Low. There were two Federally Qualified Health Centers, one rural health clinic, one ambulatory surgical center, and multiple medical groups and outpatient treatment centers throughout the region. The rate of available primary care physicians in the region was 11-13 primary care physicians per 1,000 AHCCCS children compared to the statewide rate of 23-24 per 1,000 AHCCCS children. For primary care physicians accepting AHCCCS patients, the regional rate was 9-10 physicians per 1,000 AHCCCS children. For dentists accepting AHCCCS patients, the regional rate was 6-7 dentists per 1,000 AHCCCS children compared to 16-17 dentists per 1,000 AHCCCS children statewide. The region also had three pharmacies per 1,000 AHCCCS children.

We compared the distance that regional and statewide AHCCCS children needed to travel to the nearest provider type to assist in determining whether the population in the region may have access to care issues based on travel distance. To visit the nearest primary care physician or pharmacy, 56-67% of AHCCCS children in the region traveled up to five miles services compared to 90-91% of AHCCCS children statewide. The nearest dentist and behavioral health provider were up to five miles away for 73-79% of regional AHCCCS children versus 91-92% of AHCCCS children statewide. To visit the nearest hospital 46-47% of regional AHCCCS children traveled up to 10 miles and 54% traveled more than 10 miles compared to 80-81% of AHCCCS children statewide who traveled up to five miles.

Primary Care and Well-Child Visits

Access to primary care is important for the health and well-being of children. Primary care practitioners (PCPs) provide appropriate screenings, treatment and preventive services. When children regularly visit a PCP, they are less likely to visit the emergency department for non-urgent care. Well-child visits are PCP visits scheduled at designated age intervals where a child's growth and development are measured and

⁴ Tribal affiliation refers to whether an individual is a member of a federally recognized Arizona tribe and is displayed as a flag (Yes/No) in this report. This information is captured during enrollment in AHCCCS.

tracked according to national guidelines. PCPs examine a child holistically for physical, mental, emotional and social/environmental health during a well-child visit.

Regionally, 77-78% of AHCCCS children had at least one PCP visit compared to 85-86% of AHCCCS children statewide and 86-87% of Medicaid children nationally. The regional rates did not meet the AHCCCS Minimum Performance Standard (MPS)⁵ of 84%. Two subregions met or exceeded the MPS for annual PCP visits: Show Low (all years) and Vernon (2017 and 2018). AHCCCS children ages 1-2 (82-84%) were more likely to have annual PCP visits than ages 3-5 (73-74%). A small variation was reported between Hispanic and Latino AHCCCS children (81-84%) versus Non-Hispanic or Latino AHCCCS children (75-77%) in visits to a PCP.

Regionally, 39-49% of AHCCCS children birth to 15 months had six or more well-child visits compared to 63-66% of Medicaid children nationally. The region and state rates were below the AHCCCS MPS of 65% (2017 and 2018) and 62% (2019) for this indicator. Show Low exceeded the AHCCCS statewide rate and MPS in 2019 for this indicator. For AHCCCS children ages 3-5, 48-50% of regional children had an annual well-child visit compared to 62-65% of statewide children and 72-74% of Medicaid children ages 3-6 nationally; the region was below the AHCCCS MPS of 66% for this indicator. No region met the AHCCCS statewide rates on MPS for this indicator. Hispanic or Latino AHCCCS children (50-59%) were more likely to have an annual well-child visit compared to Non-Hispanic or Latino AHCCCS children (47-49%).

Screening for Lead Poisoning

Lead poisoning is a silent killer because often there are no symptoms. Exposure to lead can cause irreversible damage to the brain and other vital organs in children, as well as intellectual and behavioral deficits. To detect abnormal blood lead levels in children, screenings are conducted via a blood lead test. According to the Arizona Department of Health Services (ADHS), children who live in areas designated as high-risk for lead poisoning should receive a blood lead test at 12 and 24 months of age, and older children who have not been previously tested should receive the blood lead test.⁶

For AHCCCS children being screened for lead poisoning one or more times by their second birthday, the regional rates increased from 17% (2017-2018) to 22% (2019) compared to AHCCCS statewide rates which increased from 32% in 2017 to 35% in 2019. Blue Ridge subregion exceeded the AHCCCS statewide rates for this indicator in 2019. Hispanic or Latino AHCCCS children (51-63%) were more likely to be screened for lead poisoning by their second birthday than Non-Hispanic or Latino AHCCCS children (33-48%).

⁶ https://www.azdhs.gov/preparedness/epidemiology-disease-control/lead-poisoning/index.php#high-risk-zip-codes-home

⁵ Minimum Performance Standard (MPS) is the minimal expected level of performance by AHCCCS Contractors. AHCCCS-reported rates are the official rates used to determine Contractor compliance with performance requirements. If a Contractor does not achieve the MPS, they will be required to submit a corrective action plan and may be subject to sanctions for each deficient measure.

Weight Assessment and Counseling⁷

Childhood obesity has both short-term and long-term effects, so it is important for PCPs to monitor weight problems in children and provide guidance for maintaining a healthy weight and lifestyle. The prevalence of obesity among children aged 2–5 years in 2015-2016 was 14% according to a national survey. For this report, we focused on AHCCCS children ages 3-5.

The regional rates for weight assessment and counseling showed AHCCCS children in Navajo/Apache were assessed for weight at 3-8% compared to AHCCCS children statewide who were assessed at rates of 9-19%. Rates for nutrition counseling were 1-2% at the regional level versus 4-5% at the state level for AHCCCS children. Physical activity assessments in the region were suppressed due to counts <6 while AHCCCS children statewide were assessed <1-1%.

Developmental Screening and Delay

During early childhood, children grow and develop at a rapid pace physically and cognitively. Although children develop skills at different times, there are guidelines that define the period when an average child should meet certain developmental milestones. National pediatric guidelines recommend developmental screenings during well-child visits for all children ages 9 months, 18 months, 2 years and 2.5 years. Developmental delay occurs when a child does not demonstrate mastery of developmental milestones. Developmental delays have been found to occur in 10-15% of preschool children nationwide.

Rates of developmental screenings in AHCCCS children birth to age 5 increased over the report period although still well below the AHCCCS median rates of 42% in 2017 and 33% in 2018. At the regional level, rates increased from 3-5% compared to statewide AHCCCS rates of 10-14%. AHCCCS children ages 1-2 (6-10%) were more likely to receive a developmental screening than age 0 (2-4%) and ages 3-5 (1-4%). Rates of diagnosing developmental delay in AHCCCS children were 2-4% at the regional level compared to 3-5% at the state level for AHCCCS children. Several subregions met or exceeded the statewide rates for diagnosing developmental delay: Blue Ridge (2017 and 2018), Concho (all years) and Show Low (all years). Male AHCCCS children (4-5%) were more likely to be diagnosed with developmental delay at the regional level than females (1-3%). AHCCCS children ages 3-5 (3-5%) were more likely to be diagnosed with developmental delay than ages 1-2 (2-4%). Of those AHCCCS children who were diagnosed with developmental delay, 29-44% of regional AHCCCS children received behavioral health services compared to 47-58% of AHCCCS children statewide. Concho exceeded the AHCCCS statewide rate for AHCCCS children receiving behavioral health services after being diagnosed with developmental delay.

⁸ Physical Activity Counseling includes sports physicals which are not provided to children in the early childhood age group.

⁷ There was limited reporting in claims data as this information was most likely collected in the medical record, so these rates should be interpreted with caution.

⁹ Due to the limited capture of developmental screenings in claims data alone, these rates should be interpreted with caution.

Behavioral Health

The social-emotional development and adaptive functioning of a young child is as important as their physical health. Negative early childhood events can lead to behavioral and physical health problems in adulthood if behavioral health intervention services are not provided at the infant and toddler stages. For young children, behavioral health services ¹⁰ would likely include day programs, crisis services, rehabilitation services, health promotion, mental health counseling, psychiatric and psychologist services, and various support services. Seven to ten percent of AHCCCS children in Navajo/Apache region received behavioral health services compared to 11-16% of AHCCCS children statewide. Male AHCCCS children in the region were more likely to receive behavioral health services (8-13%) compared to female AHCCCS children (5-7%).

Vision¹¹

Visual impairment affects a child's development, performance, and quality of life. Fortunately, most vision problems are successfully treated when detected early through regular visits to PCPs, and well-child visits should include a vision screening. It has been estimated that 20% of preschool children in the United States have eye or vision problems. Arizona's Eyes on Learning Vision Coalition recommends a vision screening beginning at age one. Children ages 3-5 should have at least one vision screening by a PCP or trained screener, and annual screenings should be provided to children in kindergarten through fourth grade.

In Navajo/Apache Region, 39-40% of AHCCCS children received an annual vision screening or well-child visit compared to 43-47% of AHCCCS children statewide. Several subregions met or exceeded the AHCCCS statewide rates for annual vision screening or well-child visit: Blue Ridge in 2017 and 2018, Concho in 2017, Show Low in 2017 and 2018 and Snowflake in all years. Annual vision screening or well-child visit rates for AHCCCS children ages 1-2 (56-57%) were higher than ages 3-5 (44-47%). Eye exams were conducted much less frequently; 5% annually at the regional level and 4-5% at the state level for AHCCCS children. Regional AHCCCS children ages 3-5 (9%) were more likely to receive an eye exam than ages 1-2 (2-3%). Follow-up eye exams were conducted on AHCCCS children in the region and statewide at rates of approximately 4-6%. AHCCCS children with visually significant eye conditions received treatment at rates of 47-60% regionally compared to 54-60% statewide.

_

¹⁰ For more detail on AHCCCS behavioral health services, visit https://www.azahcccs.gov/Members/AlreadyCovered/coveredservices.html

¹¹ Per the AHCCCS Medical Policy Manual, AHCCCS children should receive hearing and vision screenings during their well-child visits according to the periodicity schedule. Claims data does not specify each service provided during a well-child visit; thus, we cannot verify whether these screenings were provided according to the schedule. The rates in this report should be interpreted with caution.

Hearing¹¹

Most children begin hearing sounds at birth and learn to speak over time by imitating the sounds around them. However, it is reported that around two or three out of every 1,000 children are born deaf or hard-of-hearing in the United States, and more lose their hearing later in childhood. For children diagnosed with hearing loss, early detection, intervention and treatment would provide each child with the opportunity to develop better language and communication skills. Arizona strives to screen all infants before one month of age. Infants who do not pass the initial hearing screen and a rescreening, should be evaluated further to confirm or diagnose hearing loss before 3 months of age. Infants diagnosed with permanent hearing loss should receive intervention services before 6 months of age.

Around 99% (82,035) of all Arizona infants received a newborn hearing screening in 2017, which was slightly higher than the national rate of 98%. Less than 1% of all Arizona infants were diagnosed with permanent hearing loss, and of those, 42% were diagnosed before three months of age. Nationally, 10% of infants were diagnosed with permanent hearing loss, and of those, approximately 74% were diagnosed before three months of age. Additional audiology services were provided to 2-6% of AHCCCS children under age one in the region compared to 9-12% of AHCCCS children statewide.

Hearing screenings were provided to 9-12% of AHCCCS children ages 1-5 in the region compared to 20-28% of AHCCCS children statewide. Show Low exceeded the state rate for hearing screenings in AHCCCS children ages 1-5 in 2018. Male regional AHCCCS children (11-16%) were more likely to receive a hearing screening than females (7-8%). For AHCCCS children ages 1-5 who received a hearing screening, the provision of additional audiology services decreased from 94% of AHCCCS children in the region in 2017 to 87% in 2019 while statewide AHCCCS children's rates decreased from 68% to 57% over the same period. AHCCCS children ages 3-5 (97-100%) were more likely to receive additional audiology services than ages 1-2 (73-91%) in 2019. Male AHCCCS children (93-97%) were more likely to receive additional audiology services than females (80-91%).

Oral Health

Oral health is a key indicator of overall health, well-being and quality of life. Access to dental care is necessary to maintain good oral health. Two preventative care dental visits are recommended annually for children. For young children, the application of fluoride varnish to primary and permanent teeth is also recommended to prevent cavities.

In Navajo/Apache Region, 39-43% of AHCCCS children had at least one annual dental visit compared to 51-53% of AHCCCS children statewide. Neither the region nor the state met the AHCCCS MPS of 60% for annual dental visits for ages 2-20. No subregion met the AHCCCS MPS. AHCCCS children ages 3-5 (48-54%) were more likely to have at least one annual dental visit than ages 1-2 (24-27%); Hispanic or Latino AHCCCS children (41-50%) were more likely to have at least one annual dental visit than Non-Hispanic or Latino AHCCCS children (39-42%). Two preventative care dental visits are recommended annually for children.

¹¹ ibid

Regionally, 12-14% of AHCCCS children received the biannual preventative care dental visit compared to 18-20% of AHCCCS children statewide; 36-41% of regional AHCCCS children had at least one preventative care dental visit per year. Fluoride varnish was applied to 34-38% of AHCCCS children in the region compared to 47-49% of AHCCCS children statewide. AHCCCS children ages 3-5 (42-48%) were more likely to have a fluoride varnish application than ages 1-2 (21-23%). Hispanic or Latino AHCCCS children (36-46%) were more likely to have a fluoride varnish application than Non-Hispanic or Latino AHCCCS children (34-37%).

Maternal Prenatal and Postpartum Care

The health of women before pregnancy and after delivery significantly impacts their health and the health of their babies. Thus, it is important to focus on women's prenatal and postpartum care. Prenatal care involves regular visits to a health care provider to monitor the mother's health and health of the developing fetus. Women should have at least one prenatal visit in the first trimester of pregnancy. The period of up to 60 days following childbirth is called the postpartum period. Preexisting health conditions, social determinants, and newly developed conditions contribute to maternal morbidity and mortality during this period so at least one postpartum visit is recommended.

In Navajo/Apache Region, 92-95% of pregnant women began prenatal care in the first trimester compared to 84-86% of AHCCCS women statewide, which are both above the Healthy People 2030 target rate of 81%¹². All subregions met or exceeded the Healthy People 2030 target rate for timely prenatal care for all years, except for Holbrook which exceeded the target rate in 2019 only. There was little difference between AHCCCS women's age, tribal affiliation, race or ethnicity in receiving timely prenatal care as all had high rates for this indicator. For postpartum care, 96% of regional AHCCCS women had at least one postpartum visit compared to 88-89% of AHCCCS women statewide and 64-75% of Medicaid women nationally. All subregions exceeded the state rates for postpartum care in all years, except for Holbrook which exceeded the state rate in 2019 only.

Health Plan Performance

This section provided a selection of health indicators to compare results among AHCCCS children enrolled in each of the AHCCCS health plans available in the region. The following regional health plans met or exceeded the AHCCCS statewide aggregate performance and MPS for PCP visits for ages 1-5: AZ Complete Health (2017 and 2018), Banner University Family Care (2019), Comprehensive Medical and Dental Program (CMDP) (2018 and 2019), Mercy Care Plan (2018 and 2019) and UnitedHealthcare (2017 and 2019). For at least one well-child visit in the first 15 months, all regional health plans performed well (86-100%), except for the American Indian Health Program. For six or more well-child visits in the first 15 month, only CMDP exceeded the AHCCCS statewide aggregate performance and MPS in 2018. For one or more well-child visits for ages 3-5, four health plans met or exceeded the AHCCCS statewide aggregate

¹² Healthy People 2030 Prenatal Care Objective - https://health.gov/healthypeople/objectives-and-data/browse-objectives/pregnancy-and-childbirth/increase-proportion-pregnant-women-who-receive-early-and-adequate-prenatal-care-mich-08

performance and MPS: Banner University Family Care (2019), Children's Rehabilitative Services (2018), Mercy Care Plan (2018) and UnitedHealthcare (2019). Only Banner University Family Care (2019) met or exceeded the AHCCCS statewide aggregate health plan performance or MPS for preventative care dental visits for ages 1-5.¹³

Conclusion

From 2017 to 2019, Navajo/Apache showed strong performance on the following AHCCCS children's and women's health indicators: newborn hearing screenings and prenatal and postpartum care. These achievements contributed to good health outcomes throughout the region. The areas where needs were identified for AHCCCS women and children included PCP and well-child visits, the supply of health professionals, developmental screening, lead poisoning screening, vision screening and oral health. The information in this report can be combined with other available information to create a more comprehensive view of young children and women in the region for regional council planning.

¹³ The AHCCCS statewide indicator for preventative care dental visits includes ages 2-20 which incorporates a significantly larger number of AHCCCS children than our reporting on ages birth to 5, so the rates should be compared with caution.

INTRODUCTION

THE IMPORTANCE OF EARLY CHILDHOOD HEALTH

Under the direction of First Things First (FTF), the Arizona State University Center for Health Information & Research (CHiR) conducted a regional analysis of children from birth to age 5 to explore the health assets and needs in the FTF NAVAJO/APACHE Region. This report provides detailed health utilization and access to services for children birth through 5, along with prenatal and postpartum women, who were enrolled in the Arizona Health Care Cost Containment System (AHCCCS). Additional information is provided via medical board licensing data to further describe access to medical professionals and services contracted under AHCCCS. The goal is for the FTF Regional Partnership Councils to utilize the findings in this report when conducting regional planning discussions, as an additional resource and tool to the Regional Needs and Assets base report.

DEFINITIONS

Access to Care

This term refers to making health care services readily available when needed and removing all barriers.

Age Groups

- Age is defined as the age of the patient on Dec. 31 of each given year. The age for a patient is
 constant through the report year. Special age breakdowns are listed for certain indicators that
 have an associated milestone.
- Infants: less than 1 year of age
- Toddlers: greater than or equal to 1 year of age to less than 3 years of age
- Preschooler: greater than or equal to 3 years of age to less than 6 years of age

Assets

An asset is a finding where young children or women appear to be faring well regarding utilization of or access to health care.

Behavioral Health

To determine whether children are receiving behavioral health services, we used the following definition: category of service on claim equals mental health services (category of service = 47) or primary diagnosis is a behavioral health diagnosis as listed in the AHCCCS Behavioral Health Services Matrix https://www.azahcccs.gov/PlansProviders/MedicalCodingResources.html.

Children

Unless noted otherwise, all references to children denote children, ages birth to 5, who are AHCCCS members.

CMS Median

The Centers for Medicare & Medicaid Services (CMS) annually collects and reports state performance rates on a standardized set of care quality measures for Medicaid and Children's Health Insurance Program beneficiaries, called the Child and Adult Core Set. The CMS Median is the average performance among reporting states for each measure.

Habilitation

Training in independent living skills or special developmental skills, sensory-motor development, orientation and mobility and behavior intervention.

Healthcare Effectiveness Data and Information Set

The Healthcare Effectiveness Data and Information Set (HEDIS) is a tool produced by the National Committee for Quality Assurance (NCQA) that is used by most U.S. health plans to measure performance and quality in health care. HEDIS® contains over 90 measures under six domains of care: effectiveness of care, access/availability of care, experience of care, utilization and risk adjusted utilization, health plan descriptive information, and measures reported using electronic clinical data systems. The national committee collects HEDIS® survey results from health plans and Preferred Provider Organizations through the Healthcare Organization Questionnaire and collects non-survey data through the Interactive Data Submission System. HEDIS measures are specifically defined to make comparisons among health plans. The measurement set is reviewed annually. CHiR uses the AHCCCS claims within HEDIS. HEDIS measures have complicated numerator and denominator calculations, and therefore, are expressed and interpreted as rates.

Health Plans

Health plan categories include acute care, Children's Rehabilitative Services, Comprehensive Medical and Dental Program, Developmental Disability/Department of Economic Security, Long Term Care, and Fee-For-Service American Indian health plans.

Minimum Performance Standard

Minimum Performance Standard (MPS) is the minimal expected level of performance by AHCCCS Contractors. AHCCCS-reported rates are the official rates used to determine Contractor compliance with performance requirements. If a Contractor does not achieve the MPS, they will be required to submit a corrective action plan and may be subject to sanctions for each deficient measure.

Needs

A need is an area where it appears that access or utilization of health care is low.

Postpartum Period

The AHCCCS postpartum period begins the day the pregnancy terminates and continues for 60 days following pregnancy termination.

Primary Care Physician Specialties

Physicians included in the primary care specialty include Family Practitioner, General Practitioner, Internal Medicine and Pediatrician.

Race/Ethnicity

Race and ethnicity are grouped and reported in the following manner.

- Race
 - Asian/Pacific Islander
 - o Black
 - Caucasian/White
 - o Native American
 - Other/Unknown

- Ethnicity
 - Hispanic or Latino
 - Not Hispanic or Latino
 - Unknown

Up until 2017, AHCCCS only collected one race/ethnicity variable and used the Hispanic value to denote Hispanic or Latino origin. As of 2017, AHCCCS began collecting race and ethnicity as separate variables. Hispanic is retained as a race variable, but AHCCCS is phasing out its use; therefore, the decrease in the use of Hispanic in the race variable correlates to the increase in reporting of Unknown in the race variable. Ethnicity is reported separately beginning in 2018 and notes on its use in this report are below.

- Individuals who reported "Not Hispanic, Latino, Spanish" are not Hispanic or Latino origin.
- To denote those of Hispanic or Latino origin, we combine Mexican, Mexican American, Chicano, Puerto Rican, Cuban, Other Hispanic/Latino Origin, and Hispanic or Latino Unknown.
- All individuals who reported a race/ethnicity of Hispanic prior to 2017 were assigned a race of other/unknown and an ethnicity of Hispanic or Latino origin
- Ethnicity Unspecified refers to individuals who did not answer or were not provided the opportunity to give this information.
- Ethnicity Unknown means the individual chose to be unknown. AHCCCS started phasing out this category in October 2018.
- Data on multiracial individuals is not collected.

Tribal Affiliation

Tribal affiliation refers to whether an individual is a member of a federally recognized Arizona tribe and is displayed as a flag (Yes/No) in this report. This information is captured during enrollment in AHCCCS. This is based on AHCCCS-stated affiliation, not residential location.

Well-Child Visits

Children enrolled in AHCCCS receive well-child visits under the Early and Periodic Screening, Diagnostic and Treatment (EPSDT) Program. The EPSDT program provides comprehensive treatment and preventive health care services for children under age 21. The services include dental, physical, behavioral health, developmental, vision, hearing, screenings and other specialty services. EPSDT visits are all-inclusive, meaning one payment is made for all services rendered during the visit. Only certain services are billed

separately when conducted by qualified health care providers, and those are: nutritional assessments, developmental screenings, immunizations, fluoride varnish and ocular photo screening.

Women

Unless noted otherwise, all references to women denote women who were AHCCCS members.

APPROACH

CHiR and representatives from the FTF Regions, Programs, and Evaluation teams determined priority indicators for this report. FTF provided the regional and subregional boundaries. Navajo/Apache consisted of the following subregions: Blue Ridge, Concho, Holbrook, Show Low, Snowflake and Vernon. The main data source was claims data from the Arizona Health Care Cost Containment System (AHCCCS), Arizona's Medicaid agency; therefore, the results presented in this report were for children and mothers who were enrolled in AHCCCS from 2017 to 2019. This population was denoted as AHCCCS children or AHCCCS women.

AHCCCS children's health was measured in the following categories: primary care, well-child visits, health care workforce, screening for lead poisoning, weight assessment and counseling, developmental screening and delay, behavioral health, vision, hearing, oral health, immunizations, maternal prenatal and postpartum care, and health plan performance.

Many of the reported indicators were from the Healthcare Effectiveness Data and Information Set (HEDIS). HEDIS is a performance improvement tool whereby health plans, health care organizations and government agencies submit data on specific health measures. HEDIS used the collected data to calculate national performance statistics and benchmarks and set standards for measures. HEDIS specifications were applied to the AHCCCS population for each region. The denominators were listed within each indicator and are available on the National Committee for Quality Assurance website at https://www.ncqa.org/hedis/. Inclusion generally required a child to have continuous enrollment for the reporting year with no more than one gap smaller than 45 days allowed. Some indicators also required enrollment in a period preceding the reporting year.

Non-HEDIS denominators were derived from the children who met the AHCCCS inclusion criteria for the region. The AHCCCS inclusion criteria were children ages 0-5 (0 <= age < 6) who were enrolled in AHCCCS in 2017, 2018, or 2019 and residing in Arizona regions defined by First Things First. Health claims were for paid services in 2017, 2018, or 2019. Additional AHCCCS enrollment requirements were indicator-based. The complete population of children covered by AHCCCS were not included due to the limitations on AHCCCS enrollment gaps which were not met by all children.

For the distance analysis that was reported in the health care workforce section, all AHCCCS-enrolled children were assigned coordinates on a map related to their residential address on file. Health providers were also assigned coordinates from their address on file or public address, if available. Each child's address was analyzed to determine the distance in miles to the closest provider for each provider type. The children were then grouped into distance ranges as percentages. The region and state percentages

were listed side-by-side to compare totals and determine if the population in the region may have access to care issues due to the distance required to travel for health services.

Data used in this report covered all AHCCCS members in Arizona, including members living in FTF tribal regions and subregions. Report creations for tribal regions and subregions was carried out with specific approval from the tribe. For those tribal regions who did not give approval, data was included only in aggregate totals for Arizona. In the case of a tribal subregion, only aggregate totals for Arizona and the region were included.

REPORTING

There were 13 health topics discussed in this report. Each section began with context on the importance of the health topic before discussing the results from the AHCCCS claims data.

The AHCCCS results were presented at the regional level with state and national benchmarks provided for comparison, where available. When possible, the results are grouped by 1) indicators that met or were above the state average or national HEDIS standards and 2) indicators that did not meet or were below the state average or national HEDIS standards. Other notable findings were also presented that do not have comparison data. Most results were presented as percentages for standardization purposes and ease of comparison with benchmarks. The terms rate and percent were used interchangeably.

After reporting the general regional demographics, the results were displayed by gender, age group, race, ethnicity, tribal affiliation, provider type, and/or health plan when the data was available and within the data suppression guidelines stated below. Each section contained maps to display the results at the subregional level. The maps had a color gradient which compared the performance among the subregions for each indicator. A darker color denoted a higher percentage of individuals in the subregion who were included in the indicator. Percentages over 1% were rounded to the nearest whole number. Percentages less than 1% were denoted as "<1%".

A brief conclusion summarized how well the region was doing with regards to access and utilization of health care services and provided areas where the regional councils may want to focus during their regional planning conversations.

The Executive Summary was designed to provide the main findings and takeaways for the report. A definitions section explained the lesser-known terms. The data sources were detailed in the Appendix which follows the references. The report was hyperlinked for ease of navigating from the Table of Contents and the text to the associated topics, figures and tables.

To protect the confidentiality of program participants, the First Things First Data Dissemination and Suppression Guidelines preclude our reporting data related to health or developmental delay if the count is less than six. Throughout this report, information which was not available because of suppression guidelines will be indicated by entries of "<6" for counts or "DS" (data suppressed) for percentages. Data were sometimes not available for particular regions, either because a particular program did not operate in the region or because data were only available at a higher level (i.e., county, state, etc.). Cases where data were not available will be indicated by an entry of "N/A."


DATA LIMITATIONS

Most of the results in this report used AHCCCS claims and encounter data. While being limited to the population of children enrolled in Arizona Medicaid, this data source was also subject to coding errors and missing data for some indicators.

To best capture the full picture of childhood immunizations, a combination of data from claims, electronic health records, paper medical records and registry data was needed. AHCCCS used a combination of AHCCCS claims, Arizona State Immunization Information System (ASIIS) registry data and medical record data from its contractors to measure immunization rates internally. For this report, we used only AHCCCS claims as we did not have access to the other data sources. Since the AHCCCS claims data only included a subset of the immunizations of Arizona's children, our results showed substantially lower immunization rates than AHCCCS officially reports.

Per the AHCCCS Medical Policy Manual, AHCCCS children should receive hearing and vision screenings during their well-child visits according to the periodicity schedule. Claims data does not specify each service provided during a well-child visit; thus, we cannot verify whether these screenings were provided according to the schedule. The rates in this report should be interpreted with caution.

NAVAJO/APAHCE REGIONAL RESULTS



POPULATION AND DEMOGRAPHICS OF CHILDREN ENROLLED IN AHCCCS

AHCCCS exists to make care affordable to the individuals and families it enrolls, including the approximately 235,000 children birth to age 5 who were enrolled in AHCCCS from 2017 to 2019.

There were 234,616 children from birth to age 5 enrolled in AHCCCS statewide from 2017 to 2019.

(AHCCCS Claims Data, 2021)

In Navajo/Apache Region, there were 3,077 children enrolled in AHCCCS in 2017, 2,968 children enrolled in 2018 and 2,920 children enrolled in 2019. Of these, male AHCCCS children outnumbered females by 2-4% in Table 1. In Table 2, there were 6% more infants and toddlers than preschoolers in 2017, with the difference dropping to 1% more in 2019.

Approximately 65% of the AHCCCS children in Navajo/Apache Region lived in three subregions: Holbrook, Show Low and Snowflake (Figure 1). By race, 62-66% of AHCCCS children were reported as Caucasian/White, 20% were Native American, 2-3% were Black and 1% were Asian/Pacific Islander (Figure 2). By ethnicity, 82-83% were reported as Non-Hispanic or Latino and 17-18% were Hispanic or Latino (Figure 3). Affiliation with a tribal community was reported by 23-24% of AHCCCS children annually in Figure 4.¹⁴ In Figure 5, most AHCCCS children in the region were enrolled in three health plans: Steward Health Choice AZ (48-49%), UnitedHealthcare (21-31%) and the American Indian Health Plan (10%). By health claim, 29-32% were submitted by physicians, 13-15% were submitted by pharmacies and 12-13% were submitted by hospitals in Table 3.

Table 1. Number of AHCCCS Children Birth to Age 5 by Year and Sex

Year	Female	Male	Total of AHCCCS-Enrolled Children
2017	1,509	1,568	3,077
2018	1,447	1,521	2,968
2019	1,400	1,520	2,920

Source: AHCCCS Claims Data, 2021. CHIR is the source for all processing of the AHCCCS data.

Table 2. Number of AHCCCS Children Birth to Age 5 by Year and Age Group

Year	Infant (under 1)	Toddler (1-2)	Preschooler (3-5)
2017	609	1,019	1,449
2018	587	899	1,482
2019	620	827	1,473

¹⁴ Tribal affiliation refers to whether an individual is a member of a federally recognized Arizona tribe and is displayed as a flag (Yes/No) in this report. This information is captured during enrollment in AHCCCS.




Figure 1. Number of AHCCCS Children Birth to Age 5 by Subregion by Year




Figure 3. Number of AHCCCS Children Birth to Age 5 by Year and Ethnicity




Figure 4. Number of AHCCCS Children Birth to Age 5 by Year and Tribal Affiliation




Figure 5. Number of AHCCCS Children by Year and Health Plan




Table 3. Number of AHCCCS Claims by Provider Type (Billing Entity), 2017-2019

	20	17	20	18	2019		
Provider Type	Claims Count	Percent of Total	Claims Count	Percent of Total	Claims Count	Percent of Total	
Behavioral Health Outpatient Clinic	1,510	4%	900	3%	581	2%	
Dentist	2,011	6%	1,820	5%	1,900	5%	
Durable Medical Equipment Supplier	572	2%	593	2%	917	3%	
Federally Qualified Health Center (FQHC)	1,333	4%	1,016	3%	1,039	3%	
Habilitation Provider*	1,237	4%	1,108	3%	1,067	3%	
Home Health Agency	472	1%	478	1%	538	2%	
Hospital	4,359	13%	4,175	12%	4,536	13%	
Integrated Clinics**	240	1%	266	1%	475	1%	
Laboratory	824	2%	926	3%	891	2%	
Non-Emergency Transportation Providers	725	2%	849	3%	799	2%	
Occupational Therapist	231	1%	378	1%	206	1%	
Pharmacy	5,157	15%	4,642	14%	4,748	13%	
Physical Therapist	140	0%	310	1%	183	1%	
Physician – MD/DO	10,044	29%	10,617	31%	11,598	32%	
Physician Assistant	1,756	5%	1,738	5%	1,869	5%	
Registered Nurse Practitioner	2,188	6%	1,896	6%	2,233	6%	
Speech Language Pathology	115	0%	125	0%	131	0%	
Speech/Hearing Therapist	290	1%	361	1%	222	1%	
Other	1,599	5%	1,637	5%	1,908	5%	

Notes: * Habilitation is training in independent living skills or special developmental skills, sensory-motor development, orientation and mobility and behavior intervention.

^{**} An Integrated Clinic is a provider licensed by the Arizona Department of Health Services as an Outpatient Treatment Center which provides both behavioral health services and physical health services.

HEALTH CARE WORKFORCE

Currently, Arizona has 160 hospitals individually licensed by the state which are subtyped as children, critical access, long term, short term, psychiatric, rehabilitation, transplant and non-participating (Arizona Department of Health Services, 2021). The Navajo/Apache Region had one short-term hospital, Summit Healthcare Regional Medical Center in Show Low. There were two Federally Qualified Health Centers, one rural health clinic, one ambulatory surgical center, and multiple medical groups and outpatient treatment centers throughout the region (Arizona Department of Health Services, 2021).

The rate of available primary care physicians in the region was 11-13 primary care physicians per 1,000 AHCCCS children (Table 5) compared to the statewide rate of 23-24 per 1,000 AHCCCS children (Table 4). For primary care physicians accepting AHCCCS patients, the regional rate was 9-10 physicians per 1,000 AHCCCS children. For dentists accepting AHCCCS patients, the regional rate was 6-7 dentists per 1,000 AHCCCS children compared to 16-17 dentists per 1,000 AHCCCS children statewide. The region also had three pharmacies per 1,000 AHCCCS children.

In Table 6, we compared the distance that regional and statewide AHCCCS children needed to travel to the nearest provider type to assist in determining whether the population in the region may have access to care issues based on travel distance. To visit the nearest primary care physician or pharmacy, 56-67% of AHCCCS children in the region traveled up to five miles services compared to 90-91% of AHCCCS children statewide. The nearest dentist and behavioral health provider were up to five miles away for 73-79% of regional AHCCCS children versus 91-92% of AHCCCS children statewide. To visit the nearest hospital 46-47% of regional AHCCCS children traveled up to 10 miles and 54% traveled more than 10 miles compared to 80-81% of AHCCCS children statewide who traveled up to five miles.

Table 4. Supply of Key Health Professionals in Arizona per 1,000 AHCCCS Children, 2017-2019

Provider Type	20:	17	20:	18	2019*	
Provider Type	Number	Rate	Number	Rate	Number	Rate
Total Active Physicians	16,345	70	17,356	74	N/A	N/A
Active Primary Care Physicians ¹	5,396	23	5,598	24	N/A	N/A
Pediatricians ⁴	1,214	5	1,257	5	1,293	6
Active Registered and Practical Nurses ²	N/A	N/A	101,599	433	104,434	445
Dentists ³	3,796	16	3,903	17	4,012	17

Source: ¹ (Association of American Medical Colleges, 2017). (Association of American Medical Colleges, 2019). ² (National Council of State Boards of Nursing, 2021). ³ (American Dental Association, 2021). ⁴ (American Board of Pediatrics, 2020); (American Board of Pediatrics, 2019); (American Board of Pediatrics, 2018).

Notes: The rate was calculated using the Arizona population of AHCCCS children birth to age 5 (N = 234,616). National data on pediatricians excluded those who were over age 70 to better control for those who may have been deceased in recent years. ¹ Data were from the 2017 and 2019 AMA Physician Masterfiles. Active physicians were federal and non-federal with an Arizona state license who worked at least 20 hours per week. *Data on active physicians was not available for 2019.

Table 5. Supply of Key Health Professionals in Navajo/Apache Region per 1,000 AHCCCS Children, 2017-2019

Provider -		17	2018		2019	
		Rate	Num	Rate	Num	Rate
Primary Care						
Primary Care – All Licensed Primary Care Physicians ²		11	35	12	38	13
Physicians accepting AHCCCS ^{1,2} – Total	28	9	30	10	27	9
Physicians accepting AHCCCS – Pediatrics		1	3	1	3	1
Physicians accepting AHCCCS – Primary Care		8	27	9	24	8
Physicians with ≥250 AHCCCS patients per year (all ages)		6	19	6	16	5
Behavioral Health – AHCCCS ¹						
Behavioral Health Physician Specialty or Allied Health Professional		13	39	13	33	11
Primary Care with Behavioral Health Services*	2	1	3	1	6	2
Other						
Dentist – AHCCCS ²	18	6	19	6	20	7
Hospital ^{1,3}	2	1	2	1	2	1
Pharmacy ^{1,4}	8	3	8	3	8	3

Source: ¹AHCCCS Claims Data, 2021. ²Arizona Medical Board and Arizona Board of Osteopathic Medical Examiners in Medicine and Surgery, 2021. ³ (Arizona Department of Health Services, 2021). ⁴RXOpen dataset, accessed from data.gov, 2020. CHiR was the source for all processing of the AHCCCS and Workforce data.

Notes: The rate was calculated using the regional population of AHCCCS children birth to age 5 in 2019 (N = 3,077 for 2017, N = 2,968 for 2018 and N = 2,920 for 2019). Pharmacies that were co-located with hospitals or clinics were not captured in the data. Hospital, and pharmacy historic data was not available, so all numbers are based on most recent data available.

^{*}This includes Federally Qualified Health Clinics and Integrated Clinics. These facilities provide both behavioral health services and physical health services.

Table 6. Percent of AHCCCS Children Grouped by Travel Distance Between Provider and Child's Residence by Provider Type for Region and Arizona, 2017-2019

Dunaidau Taura / Nailas	V	0-1 M	liles	1-5 Miles		5-10 Miles		10+ Miles		Unknown**	
Provider Type/Miles	Year	Region	ΑZ	Region	ΑZ	Region	ΑZ	Region	ΑZ	Region	ΑZ
Behavioral Health	2017	38%	62%	37%	30%	5%	3%	19%	2%	1%	4%
Specialty or Primary Care with Behavioral Health	2018	40%	64%	37%	27%	6%	3%	14%	2%	4%	3%
Services*	2019	41%	65%	32%	27%	5%	3%	19%	2%	3%	4%
	2017	31%	62%	47%	29%	7%	3%	14%	4%	1%	2%
Dentist	2018	30%	63%	44%	29%	7%	3%	19%	3%	1%	3%
	2019	34%	63%	45%	28%	6%	3%	13%	3%	1%	3%
	2017	5%	11%	30%	69%	12%	9%	54%	10%	<1%	<1%
Hospital	2018	5%	12%	30%	69%	11%	9%	54%	11%	<1%	<1%
	2019	5%	12%	30%	69%	11%	9%	54%	11%	<1%	<1%
	2017	23%	64%	39%	26%	6%	3%	32%	7%	<1%	<1%
Pharmacy	2018	24%	64%	37%	26%	7%	3%	32%	7%	<1%	<1%
	2019	24%	64%	38%	26%	6%	3%	31%	7%	<1%	<1%
Primary Care Physician	2017	26%	56%	32%	34%	6%	4%	33%	4%	4%	3%
	2018	26%	56%	30%	35%	6%	3%	34%	4%	4%	3%
	2019	30%	57%	37%	34%	8%	3%	21%	4%	4%	3%

Source: ¹ (AHCCCS Claims Data, 2021). ² (Arizona Medical Board and Arizona Board of Osteopathic Medical Examiners in Medicine and Surgery, 2021). CHiR was the source for all processing of the AHCCCS and Workforce data.

Notes: See the Approach section for details on this methodology. Pharmacies that were co-located with hospitals or clinics were not captured in the data. Historic data on Hospital and Pharmacy were not available, so all numbers are based on the most recent data available. *Behavioral Health providers includes primary care providers that offer behavioral health services. **The Unknown column captured children who did not have an exact-match address, so the number of miles to the nearest provider could not be accurately calculated.

PRIMARY CARE

Access to primary care is important for the health and well-being of children. Primary care practitioners (PCPs) provide appropriate screenings, treatment and preventive services. When children regularly visit a PCP, they are less likely to visit the emergency department for non-urgent care (Transforming Clinical Practice Initiative, 2019) (Piehl, Clemens, & Joines, 2000).

The inclusion criteria for this indicator were children enrolled in the previous 12 months who had at least one claim with a primary care provider, which includes primary care physicians, nurse practitioners and physician assistants.

Statewide, 85-86% of AHCCCS children ages 25 months to six years had at least one annual PCP visit from 2017 to 2019.

(Arizona Health Care Cost Containment System, July 2021) Regionally in Table 7, 77-78% of AHCCCS children had at least one PCP visit compared to 85-86% of AHCCCS children statewide and 86-87% of Medicaid children nationally. The regional rates did not meet the AHCCCS Minimum Performance Standard (MPS) ¹⁵ of 84%. In Figure 6, two subregions met or exceeded the MPS for annual PCP visits: Show Low (all years) and Vernon (2017 and 2018).

AHCCCS children ages 1-2 (82-84%) were more likely to have annual PCP visits than ages 3-5 (73-74%). A small variation was reported between Hispanic and Latino AHCCCS children (81-84%) versus Non-Hispanic or Latino AHCCCS children (75-77%) in visits to a PCP (Figure 7).

Table 7. Arizona and Regional AHCCCS Rates for PCP Visits, 2017-2019

Indicator/Voor	20	17	20	18	2019		
Indicator/Year	Region	Arizona	Region	Arizona	Region	Arizona	
Access to Primary Care	77%	85%	77%	85%	78%	86%	

Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

_

¹⁵ Minimum Performance Standard (MPS) is the minimal expected level of performance by AHCCCS Contractors. AHCCCS-reported rates are the official rates used to determine Contractor compliance with performance requirements. If a Contractor does not achieve the MPS, they will be required to submit a corrective action plan and may be subject to sanctions for each deficient measure.





Figure 6. Percent of AHCCCS Children Ages 1 – 5 with a Visit to PCP by Subregion by Year


Regional Range 50.8% 87.0%

Figure 7. Percent of AHCCCS Children Ages 1-5 with a Visit to PCP by Age Group, Tribal Affiliation, Ethnicity, Race and Year


Percent of children with a visit to primary care practitioner ages 1 - 5 by age group




Percent of children with a visit to primary care practitioner ages 1 - 5 by ethnicity



Percent of children with a visit to primary care practitioner ages 1 - 5 by tribal affiliation



Percent of children with a visit to primary care practitioner ages 1 - 5 by race



WELL-CHILD VISITS

Well-child visits provide an opportunity for PCPs to examine a child holistically for physical, mental, emotional and social/environmental health. A child's growth and development are tracked during a well-child visit. Screenings, counseling and immunizations take place at well-child visits. PCPs can instill healthy behaviors in children by reinforcing their importance during well-child visits. Parents and caregivers can team up with PCPs to address concerns. Creating a trusted relationship between the PCP and child is important as the child ages and develops, so these visits are beneficial to everyone involved (Moreno, 2018); (Sturgeon, 2015).

This HEDIS indicator assesses whether children who turned 15 months old during the measurement year had one or more well-child visits since birth, categorized by number of visits from one to six or more. A separate HEDIS indicator assesses whether children ages 3-5 had an annual well-child visit.


Regionally in Table 8, 39-49% of AHCCCS children birth to 15 months had six or more well-child visits compared to 63-66% of Medicaid children nationally. The region and state rates were below the AHCCCS MPS of 65% (2017 and 2018) and 62% (2019) for this indicator. In Figure 8, 84-88% of these children had at least one well-child visit. Show Low exceeded the AHCCCS statewide rate and MPS in 2019 for this indicator in Figure 9.

For AHCCCS children ages 3-5 in Table 8, 48-50% of regional children had an annual well-child visit compared to 62-65% of statewide children and 72-74% of Medicaid children ages 3-6 nationally. The region was below the AHCCCS MPS of 66% for this indicator. No region met the AHCCCS statewide rates on MPS for this indicator in Figure 11. In Figure 12, Hispanic or Latino AHCCCS children (50-59%) were more likely to have an annual well-child visit compared to Non-Hispanic or Latino AHCCCS children (47-49%).

Table 8. Arizona and Regional AHCCCS Rates for Well-Child Visits, 2017-2019

Indicator/Year	2017		2018		2019	
	Region	Arizona	Region	Arizona	Region	Arizona
Six or More Well-Child Visits in First 15 Months of Life	43%	53%	39%	58%	49%	60%
Annual Well-Child Visit, Ages 3-5	49%	62%	50%	63%	48%	65%

Figure 8. Percent of Regional AHCCCS Children by Number of Well-Child Visits Completed During Their First 15 Months by Number of Visits and Year







Figure 9. Percent of AHCCCS Children Who Had Six or More Well-Child Visits During Their First 15 Months of Life by Subregion and Year


19.3% 63.2%

Figure 10. Percent of AHCCCS Children Who Had Six or More Well-Child Visits During Their First 15 Months of Life by Race, Ethnicity, Tribal Affiliation and Year

Percent of children with six or more well-child visits during their first 15 months of life by race




Percent of children with six or more well-child visits during their first 15 months of life by ethnicity



Percent of Children

Percent of children with six or more well-child visits during their first 15 months of life by tribal affiliation



Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

Note: Data was suppressed for Asian/Pacific Islander and Black.





Figure 11. Percent of AHCCCS Children Ages 3-5 with a Well-Child Visit by Subregion by Year


34.5% 58.9%

Figure 12. Percent of AHCCCS Children Ages 3-5 with a Well-Child Visit by Race, Ethnicity, Tribal Affiliation and Year

Percent of children ages 3-5 with well-child visit by race




Percent of children ages 3-5 with well-child visit by ethnicity



Percent of children ages 3-5 with a well-child visit by tribal affiliation

Percent of Children



SCREENING FOR LEAD POISONING

Exposure to lead can cause damage to the brain and other vital organs, as well as intellectual and behavioral deficits. Because children who are exposed to lead often have no obvious symptoms, lead poisoning often goes unrecognized. Screening for lead via a capillary or venous lead blood test is an easy way to detect an abnormal blood lead level in children. There is no safe blood lead level. If not found early, exposure to lead and high blood lead levels can lead to irrevocable effects on a child's physical and mental health (Arizona Department of Health Services, 2006); (Arizona Department of Health Services, 2003); (National Center for Environmental Health, 2020).

In Arizona, blood lead results are reportable to the Arizona Department of Health Services (ADHS) for children less than six years old. According to ADHS, children who live in areas designated as high-risk¹⁶ for lead poisoning should receive a blood lead test at 12 and 24 months of age, and older children who have not been previously tested should receive a blood lead test. ADHS reported 61,391 children under age six (14% of children under age 5) were screened in 2019, and 40,773 (66%) of those children lived in high-risk areas. Of the children living in high-risk areas, 29% were screened at 12 months of age, and 19% were screened at 24 months of age. Only 10% of children were screened at both intervals (Arizona Department of Health Services, 2021).

For AHCCCS children being screened for lead poisoning one or more times by their second birthday in Table 9, the regional rates increased from 17% (2017-2018) to 22% (2019) compared to AHCCCS statewide rates which increased from 32% in 2017 to 35% in 2019. Blue Ridge subregion exceeded the AHCCCS statewide rates for this indicator in 2019 (Figure 13). In Figure 14, Hispanic or Latino AHCCCS children (51-63%) were more likely to be screened for lead poisoning by their second birthday than Non-Hispanic or Latino AHCCCS children (33-48%).

Table 9. Arizona and Regional AHCCCS Rates for Lead Poisoning Screening, 2017-2019

Indicator/Year	20	17	20	18	20	19
indicatory rear	Region	Arizona	Region	Arizona	Region	Arizona
One or More Tests for Lead Poisoning by Second Birthday	17%	32%	17%	34%	22%	35%

¹⁶ Interactive map of Arizona neighborhoods to identify those considered to be high-risk is online at http://www.azhealth.gov/leadmap




Figure 13. Percent of AHCCCS Children Who Had One or More Tests for Lead Poisoning by Their Second Birthday by Subregion by Year


11.8% 36.7%


Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

Note: Data was suppressed for Concho in 2019, Holbrook in 2019, Snowflake in 2018 and Vernon in 2018.


Figure 14. Percent of AHCCCS Children Who Had One or More Tests for Lead Poisoning by Their Second Birthday by Race, Ethnicity, Tribal Affiliation and Year

Percent of children who had one or more capillary or venous lead blood test for lead poisoning by their second birthday by race Percent of children who had one or more capillary or venous lead blood test for lead poisoning by their second birthday by ethnicity





Percent of children who had one or more capillary or venous lead blood test for lead poisoning by their second birthday by tribal affiliation



Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

Note: Data was suppressed for Asian Pacific Islander and Black.

WEIGHT ASSESSMENT AND COUNSELING¹⁷

Childhood obesity has both short-term and long-term effects, so it is important for PCPs to monitor weight problems in children and provide guidance for maintaining a healthy weight and lifestyle. The prevalence of obesity among children aged 2–5 years in 2015-2016 was 14% according to the National Health and Nutrition Examination Survey (Hales, Carroll, Fryar, & Ogden, 2017). For this report, we focused on AHCCCS children ages 3-5.

The regional rates for weight assessment ¹⁸ and counseling in Figure 15 showed AHCCCS children in Navajo/Apache were assessed for weight at 3-8% compared to AHCCCS children statewide in Table 10 who were assessed at rates of 9-19%. Rates for nutrition counseling were 1-2% at the regional level versus 4-5% at the state level for AHCCCS children. Physical activity assessments in the region were suppressed due to counts <6 while AHCCCS children statewide were assessed <1-1%. ¹⁹

The national HEDIS Medicaid rates were reported in Table 11; these rates included children ages 3-17, and therefore, were not strictly comparable to the region or state rates for AHCCCS children ages birth to 5.

Table 10. Arizona AHCCCS Rates for Weight Assessment and Counseling, Ages Birth to 5, 2017-2019

Indicator/Year	2017	2018	2019
BMI Assessment	9%	12%	19%
Nutrition Counseling	4%	5%	5%
Physical Activity Counseling	<1%	1%	1%

Source: AHCCCS Claims Data, 2021. CHIR is the source for all processing of the AHCCCS data.

_

¹⁷ There was limited reporting in claims data as this information was most likely collected in the medical record, so these rates should be interpreted with caution.

¹⁸ Under HEDIS, the rates for weight assessment are an evaluation of whether Body mass index (BMI) percentile is accessed and does not determine the absolute BMI value. The diagnosis codes for pediatric BMI included: Z68.51 (< 5th percentile for age), Z68.52 (5th percentile to < 85th percentile for age), Z68.53 (85th percentile to < 95th percentile for age) and Z68.54 (≥ 95th percentile for age).


¹⁹ Physical Activity Counseling includes sports physicals which are not provided to children in the early childhood age group.

Table 11. National Medicaid HEDIS Rates for Weight Assessment and Counseling, Ages 3-17 Years, 2017-2019

Indicator/Year	2017	2018	2019
BMI Assessment	73%	74%	77%
Nutrition Counseling	67%	67%	68%
Physical Activity Counseling	61%	62%	64%

Source: (National Committee for Quality Assurance, 2021).

Figure 15. Percent of AHCCCS Children Ages 3-5 Who Had Weight Assessment and Counseling by Year



Percent of Children

Source: AHCCCS Claims Data, 2021. CHIR is the source for all processing of the AHCCCS data.

Notes: Each of the three items above is a different indicator. Because BMI norms for youth vary with age and sex, this indicator evaluates whether BMI percentile is assessed, rather than an absolute BMI value.

DEVELOPMENTAL SCREENING AND DELAY

During early childhood, children grow and develop at a rapid pace physically and cognitively. Although children develop skills at different times, there are guidelines that define the period when an average child should meet certain developmental milestones. The American Academy of Pediatrics recommends developmental screenings during well-child visits for all children ages 9 months, 18 months, 2 years and 2.5 years (Centers for Disease Control and Prevention). Parents may also notice concerns they have about their child's development and discuss them with their child's health care provider.

Developmental delay occurs when a child does not demonstrate mastery of developmental milestones, and the delay can range from mild to severe. Developmental delays have been found to occur in 10-15% of preschool children (Choo, Agarwal, How, & Yeleswarapu, 2019). The National Health Interview Survey found that from 2015-2018, 18% of U.S. children ages 3-17 years had at least one developmental disability (Zablotsky & Black, 2020). After being diagnosed with a developmental delay, children should be referred to appropriate behavioral health services.

AHCCCS PCPs use developmental screening tools during 9-month, 18-month and 24-month well-child visits. Developmental screenings are assessed in claims data using billing code CPT 96110. AHCCCS has an active Performance Improvement Project to increase the number of screenings in its eligible populations (Arizona Health Care Cost Containment System, 2021). AHCCCS analyzed its own performance on developmental screenings using several data sources and reported 26% (Median = 42%) of eligible members in acute care screened in 2017 and 30% (Median = 33%) screened in 2018. Rates for AHCCCS children in foster care were 34% and 38% for the same years, respectively. AHCCCS also analyzed the 2018 data for disparities and found disparities in five of Arizona's 15 Counties: Apache, Gila, Navajo, Santa Cruz and Yavapai. Racial disparity was also demonstrated for the American Indian population.

Table 12 and Figure 16 showed rates of developmental screenings in AHCCCS children birth to age 5 increased over the report period although still well below the AHCCCS median rates. At the regional level, rates increased from 3-5% compared to statewide AHCCCS rates of 10-14%. ²⁰ In Figure 17, AHCCCS children ages 1-2 (6-10%) were more likely to receive a developmental screening than age 0 (2-4%) and ages 3-5 (1-4%).

Rates of diagnosing developmental delay in AHCCCS children were 2-4% at the regional level compared to 3-5% at the state level for AHCCCS children in Table 12. Several subregions met or exceeded the statewide rates for diagnosing developmental delay in Figure 18: Blue Ridge (2017 and 2018), Concho (all years) and Show Low (all years). In Figure 19, male AHCCCS children (4-5%) were more likely to be diagnosed with developmental delay at the regional level than females (1-3%). AHCCCS children ages 3-5 (3-5%) were more likely to be diagnosed with developmental delay than ages 1-2 (2-4%).

Of those AHCCCS children who were diagnosed with developmental delay, 29-44% of regional AHCCCS children received behavioral health services compared to 47-58% of AHCCCS children statewide (Table 12). Concho exceeded the AHCCCS statewide rate for AHCCCS children receiving behavioral health services after being diagnosed with developmental delay in Figure 20.

Table 12. Arizona and Regional AHCCCS Rates for Developmental Screenings and Delay, 2017-2019

Indicator (Vacu	20	2017		2018		19
Indicator/Year	Region	Arizona	Region	Arizona	Region	Arizona
Developmental Screening, Ages Birth to 5	3%	10%	5%	11%	5%	14%
Diagnosing Developmental Delay, Ages Birth to 5	2%	3%	3%	4%	4%	5%
Developmental Delay and Behavioral Health Services, Ages 3-5	44%	49%	29%	47%	41%	58%

Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

_


 $^{^{20}}$ Due to the limited capture of developmental screenings in claims data alone, these rates should be interpreted with caution.

Table 13. Percent of Claims by Provider Type for AHCCCS Children Receiving Developmental Screenings, 2017-2019

	2017		2018		2019	
Provider Type	Claims	Percent	Claims	Percent	Claims	Percent
Trovider Type	Count	of Total	Count	of Total	Count	of Total
Federally Qualified Health Center (FQHC)	<6	DS	8	4%	17	10%
Physician – MD/DO	74	80%	173	92%	150	85%
Registered Nurse Practitioner	12	13%	6	3%	7	4%

Notes: Not all provider categories are shown, so percentages may not sum to 100%. Data was suppressed for Physician Assistant.

Figure 16. Percent of AHCCCS Children Receiving Developmental Screenings, Age 1-2, by Subregion by Year




Regional Range 4.3% 16.4%

Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.


Note: Data was suppressed for Holbrook (2017 and 2018), Snowflake (2018) and Vernon (2017).

Figure 17. Percent of AHCCCS Children Receiving Developmental Screenings by Age Group, Tribal Affiliation, Ethnicity, Race and Year


Percent of children receiving developmental screenings over all children by age group




Percent of children receiving developmental screenings over all children ages 1-2 by tribal affiliation



Percent of children receiving developmental screenings over all children ages 1-2 by ethnicity



Percent of children receiving developmental screenings over all children ages 1-2 by race



Notes: Since developmental delay screenings are more likely to take places for those ages 1-2, the other analyses focus on that age group. Data was suppressed for Asian/Pacific Islander and Black.




Figure 18. Percent of AHCCCS Children with a Diagnosed Developmental Delay by Subregion by Year


Regional Range

5.6%


1.4%

Figure 19. Percent of AHCCCS Children with a Diagnosed Developmental Delay by Ethnicity, Race, Sex, Tribal Affiliation, Age Group and Year


Percent of children with a diagnosed developmental delay over all children by ethnicity




Percent of children with a diagnosed developmental delay over all children by race




Percent of children with a diagnosed developmental delay over all children by sex



Percent of children with a diagnosed developmental delay over all children by tribal affiliation



Percent of children with a diagnosed developmental delay over all children by age group



Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

Note: Data was suppressed for Asian/Pacific Islander, Black and age 0.

2017 2018 2019 Holbrook Holbrook Holbrook 71.4% Concho Concho 38.1% Concho 45.5% 36.4% Snowflake Snowflake 39.5% Snowflake ShowLow ShowLow ShowLow 35.3% 31.6% BlueRidge Vernon Vernon BlueRidge BlueRidge Vernon © 2022 Mapbox © OpenStreetMap Regional Range


Figure 20. Percent of AHCCCS Children with a Diagnosed Developmental Delay Who Have Received Behavioral Health Services by Subregion by Year

31.6%

71.4%


Note: Data was suppressed for Blue Ridge (2017), Concho (2017 and 2018), Holbrook (all years), Snowflake (2017 and 2018) and Vernon (all years).

Figure 21. Percent of AHCCCS Children with a Diagnosed Developmental Delay Who Have Received Behavioral Health Services by Sex and Year



Notes: The indicator uses diagnosis code R62.50: Unspecified lack of expected normal physiological development in childhood. Data was suppressed for Hispanic of Latino (2017 and 2018), female (2017), Asian/Pacific Islander (all years), Black (all years), tribally affiliated (2017 and 2018), age 0 (all years) and age 1-2 (2018).

Figure 22. Percent of AHCCCS Children with a Diagnosed Developmental Delay Who Have Received Behavioral Health Services by Age Group and Year



Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

Notes: The indicator uses diagnosis code R62.50: Unspecified lack of expected normal physiological development in childhood. Data for newborns was suppressed for all years.

Table 14. Percent of Claims by Provider Type for AHCCCS Children with a Diagnosed Developmental Delay Who Have Received Behavioral Health Services, 2017-2019

	20	17	2018		2019	
Provider Type	Claims Count	Percent of Total	Claims Count	Percent of Total	Claims Count	Percent of Total
Behavioral Health Outpatient Clinic	129	41%	108	36%	77	21%
Durable Medical Equipment Supplier	43	14%	27	9%	12	3%
Hospital	60	19%	53	18%	36	10%
Integrated Clinics	27	9%	51	17%	141	38%
Occupational Therapist	<6	DS	<6	DS	16	4%
Physician – MD/DO	16	5%	12	4%	20	5%
Speech/Hearing Therapist	6	2%	<6	DS	28	8%
Other	35	11%	51	17%	33	9%

Note: Data was suppressed for Speech Language Pathology.

BEHAVIORAL HEALTH

During the early years of life, the social-emotional development and adaptive functioning of a child changes rapidly and profoundly as their developing brains encounter experiences (National Scientific Council on the Developing Child, 2004). The Adverse Childhood Experiences studies demonstrate how negative early childhood events such as neglect, violence and trauma can lead to behavioral and physical health problems in adulthood like chronic disease, mental illness, and substance abuse (Centers for Disease Control and Prevention, n.d.). However, these effects can be mitigated with proper intervention at the infant and toddler stages by behavioral health services (Arizona Health Care Cost Containment System, 2018). For young children, behavioral health services²¹ would likely include day programs, crisis services, rehabilitation services, health promotion, mental health counseling, psychiatric and psychologist services, and various support services.

Pediatric behavioral health providers screen AHCCCS children from birth to age five for emotional, behavioral, and/or developmental needs. A national screening tool assists providers in coordinating services based on the intensity of need and formulating an integrated treatment plan (American Academy of Child and Adolescent Psychiatry, 2006).

²¹ For more detail on AHCCCS behavioral health services, visit https://www.azahcccs.gov/Members/AlreadyCovered/coveredservices.html

Of AHCCCS children statewide, 11% of children received behavioral services in 2017, nearly 12% of children in 2018 and nearly 16% of children in 2019.


According to Table 15, 7-10% of AHCCCS children in Navajo/Apache region received behavioral health services compared to 11-16% of AHCCCS children statewide. In Figure 24, male AHCCCS children in the region were more likely to receive behavioral health services (8-13%) compared to female AHCCCS children (5-7%).

Table 15. Arizona and Regional AHCCCS Rates for Behavioral Health Services, Ages 3-5, 2017-2019

Indicator/Year	2017		2018		2019	
maicatory rear	Region	Arizona	Region	Arizona	Region	Arizona
Behavioral Health Services, Ages 3-5	8%	11%	7%	12%	10%	16%

Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.


Figure 23. Percent of AHCCCS Children Ages 3-5 Receiving Behavioral Health Services by Subregion by Year




Regional Range 3.5% 15.2%

Figure 24. Percent of AHCCCS Children Ages 3-5 Receiving Behavioral Health Services by Sex, Tribal Affiliation, Ethnicity, Race and Year


Percent of children who are receiving behavioral health services of all children ages 3-5 by sex




Percent of children who are receiving behavioral health services of all children ages 3-5 by tribal affiliation



Percent of children who are receiving behavioral health services of all children ages 3-5 by ethnicity



Percent of children who are receiving behavioral health services of all children ages 3-5 by race



Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

Note: Data was suppressed for Asian/Pacific Islander (all years) and Black (2018 and 2019).

Table 16. Percent of Claims by Provider Type for AHCCCS Children Ages 3-5 Receiving Behavioral Health Services, 2017-2019

	20	17	20	18	20	19
Provider Type	Claims Count	Percent of Total	Claims Count	Percent of Total	Claims Count	Percent of Total
Behavioral Health Outpatient Clinic	1,142	66%	762	53%	491	35%
Durable Medical Equipment Supplier	58	3%	40	3%	33	2%
Habilitation Provider	<6	DS	95	7%	<6	DS
Hospital	165	10%	126	9%	128	9%
Integrated Clinics	175	10%	196	14%	419	30%
Occupational Therapist	28	2%	38	3%	44	3%
Physical Therapist	<6	DS	19	1%	37	3%
Physician – MD/DO	39	2%	28	2%	62	4%
Speech Language Pathology	<6	DS	22	2%	8	1%
Speech/Hearing Therapist	38	2%	14	1%	69	5%
Other	83	5%	104	7%	117	8%

VISION

Health conditions such as vision problems are detected through regular visits to PCPs. The American Public Health Association estimates that 20% of preschoolers have eye or vision problems (American Public Health Association, 2019). Vision screenings check the appearance of the eyes and detect potential eye problems. Most vision problems are successfully treated when detected early, but many children do not receive adequate vision screenings. A lack of vision care at younger ages can mean higher rates of undetected vision problems, leading to visual impairments that affect a child's development, performance, and quality of life.

Of all Arizona children 0-5 years old, 35% received a vision screening 2019-2020.

(Child and Adolescent Health Measurement Initiative)

Arizona's Eyes on Learning Vision Coalition recommends a vision screening as early as age one during a well-child visit. Other settings that provide vision screening include pediatrician offices,

educational settings and community settings. Children ages 3-5 should have at least one vision screening by a PCP or trained screener during this timeframe. Annual screenings should be provided to children in kindergarten through fourth grade. A vision screening is not necessary for children with certain developmental delays that cause difficulties with language and speech, motor skills, behavior, memory, learning, or other neurological functions. Instead, eye doctors recommend that all children with these types of delays receive a comprehensive eye exam. (Eyes on Learning, n.d.)

Vision screenings are typically included in AHCCCS' well-child visits according to their vision periodicity schedule and as medically necessary (Arizona Health Care Cost Containment System, 2021). However, the vision screening is not billed as a separate claim when completed during a well-child visit. Therefore, to capture the population of children who received a vision screening, we assumed that AHCCCS children were screened at their annual well-child visit, or they received a separately billable vision screening. Additional analysis showed that there were very few children who received a vision screening and not a well-child visit annually. Given that the claims data did not specify that a vision screening occurred during the well-child visit, these rates should be interpreted with caution and may be an overestimation of actual vision screenings.

Eye exams are completed by optometrists or ophthalmologists, so we captured those using procedure codes for ophthalmological services. We designated the eye exam as a follow-up eye exam if the visit occurred within six months of a vision screening or well-child visit. If a child was diagnosed with a visually significant eye condition during an eye exam and received treatment or additional visits to an optometrist or ophthalmologist for eyeglasses, surgery or other procedures, the rate of treatment was reported under "visually significant eye conditions who receive treatment". To calculate the rate for visually significant eye conditions who receive treatment, the denominator was all AHCCCS children who received an eye exam and had a diagnosis of strabismus, refraction and accommodation, amblyopia, or other eye disorders; and of those AHCCCS children with an eye condition, the numerator included children who were treated for the eye condition.

In Navajo/Apache Region, 39-40% of AHCCCS children received an annual vision screening or well-child visit compared to 43-47% of AHCCCS children statewide (Table 17). Several subregions met or exceeded the AHCCCS statewide rates for annual vision screening or well-child visit in Figure 25: Blue Ridge in 2017

and 2018, Concho in 2017, Show Low in 2017 and 2018 and Snowflake in all years. Annual vision screening or well-child visit rates for AHCCCS children ages 1-2 (56-57%) were higher than ages 3-5 (44-47%) (Figure 26).

Eye exams were conducted much less frequently; 5% annually at the regional level and 4-5% at the state level for AHCCCS children in Table 17. Regional AHCCCS children ages 3-5 (9%) were more likely to receive an eye exam than ages 1-2 (2-3%), in Figure 28.

Follow-up eye exams were conducted on AHCCCS children in the region and statewide at rates of approximately 4-6%. AHCCCS children with visually significant eye conditions received treatment at rates of 47-60% regionally compared to 54-60% statewide.

Table 17. Arizona and Regional AHCCCS Rates for Vision, 2017-2019

Indicator/Year	20	2017		2018		19
mulcatory rear	Region	Arizona	Region	Arizona	Region	Arizona
Vision Screening or Well-Child Visit	40%	44%	40%	43%	39%	47%
Eye Exams	5%	4%	5%	4%	5%	5%
Eye Exams after Vision Screening or Well-Child Visit	4%	4%	5%	5%	6%	4%
Visually Significant Eye Conditions Who Receive Treatment	50%	54%	47%	56%	60%	60%




Figure 25. Percent of AHCCCS Children Receiving Vision Screening or Well-Child Visit by Subregion by Year


Regional Range

12.7% 53.3%

Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.


Note: These results are under review and not finalized until delivery of the final report.

Figure 26. Percent of AHCCCS Children Receiving Vision Screening or Well-Child Visit by Age Group, Tribal Affiliation, Ethnicity, Race and Year







Percent of children receiving vision screening or well child visit by tribal affiliation



Percent of children receiving vision screening or well child visit by ethnicity



Percent of children receiving vision screening or well child visit by race







Figure 27. Percent of AHCCCS Children Receiving Eye Exams by Subregion by Year

Regional Range

2.5% 10.7%

Figure 28. Percent of AHCCCS Children Receiving Eye Exams by Age Group, Tribal Affiliation, Ethnicity, Race and Year



Notes: Eye exams are performed by an optometrist or ophthalmologist. Data was suppressed for Asian/Pacific Islander and Black.

HEARING

Most children begin hearing sounds at birth and learn to speak over time by imitating the sounds around them (NIDCD Information Clearinghouse, 2011). The National Institute on Deafness and Other Communication Disorders reports that around two or three out of every 1,000 children are born deaf or hard-of-hearing in the United States, and more lose their hearing later in childhood (NIDCD Information Clearinghouse, 2011). To detect hearing loss early, every state conducts universal newborn hearing screenings before a baby is discharged from a hospital or birthing center. If hearing loss is indicated, parents will be referred to an audiologist to conduct more comprehensive hearing testing and determine appropriate intervention services. For children diagnosed with hearing loss, early intervention helps children develop better language and communication skills.

Arizona strives to screen all infants before 1 month of age. Infants who do not pass the initial hearing screen and a rescreening, should be evaluated further to confirm or diagnose hearing loss before 3 months of age. Infants diagnosed with permanent hearing loss should receive intervention services before 6 months of age (Arizona Department of Health Services, n.d.). This report included available data on hearing screenings along with comprehensive hearing testing, evaluation and assessment which were termed "additional audiology services".

Around 99% (82,035) of all Arizona infants received a newborn hearing screening in 2017 (Arizona Health Care Cost Containment System, 2018) which was slightly higher than the national rate of 98% (National Center on Birth Defects and Developmental Disabilities, 2019). Less than 1% of all Arizona infants were diagnosed with permanent hearing loss, and of those, 42% were diagnosed before three months of age (Arizona Health Care Cost Containment System, 2018). Nationally, 10% of infants were diagnosed with permanent hearing loss, and of those, approximately 74% were diagnosed before three months of age (National Center on Birth Defects and Developmental Disabilities, 2019). Additional audiology services were provided to 2-6% of AHCCCS children under age one in the region compared to 9-12% of AHCCCS children statewide (Table 18).


Hearing screenings were provided to 9-12% of AHCCCS children ages 1-5 in the region compared to 20-28% of AHCCCS children statewide in Table 18. In Figure 30, Show Low exceeded the state rate for hearing screenings in AHCCCS children ages 1-5 in 2018. In Figure 31, male regional AHCCCS children (11-16%) were more likely to receive a hearing screening than females (7-8%).

For AHCCCS children ages 1-5 who received a hearing screening, the provision of additional audiology services decreased from 94% of AHCCCS children in the region in 2017 to 87% in 2019 while statewide AHCCCS children's rates decreased from 68% to 57% over the same period (Table 18). In Figure 33, AHCCCS children ages 3-5 (97-100%) were more likely to receive additional audiology services than ages 1-2 (73-91%) in 2019. Male AHCCCS children (93-97%) were more likely to receive additional audiology services than females (80-91%).

Table 18. Percent of AHCCCS Statewide and Regional Hearing Results, 2017-2019

Indicator / Year	2017		20	2018		2019	
indicator / rear	Region	Arizona	Region	Arizona	Region	Arizona	
Additional Audiology Services Under Age One	5%	11%	6%	12%	2%	9%	
Hearing Screening Ages 1-5	9%	20%	12%	22%	9%	28%	
Additional Audiology Services for those Screened, Ages 1-5	94%	68%	91%	66%	87%	57%	

Figure 29. Percent of AHCCCS Children Under 1 Year of Age Who Had Audiology Services by Sex and Year



Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

Note: Data was suppressed for Male in 2019.

Table 19. Percent of Claims by Provider Type for AHCCCS Children Ages 1-5 Receiving Hearing Screening Tests, 2017-2019

	2017		20	18	2019	
Provider Type	Claims Count	Percent of Total	Claims Count	Percent of Total	Claims Count	Percent of Total
Audiologist	46	82%	62	82%	44	77%
Physician – MD/DO	9	16%	13	17%	10	18%





Figure 30. Percent of AHCCCS Children Ages 1-5 Receiving Hearing Screening Tests by Subregion by Year


Regional Range
2.6% 27.3%

Figure 31. Percent of AHCCCS Children Ages 1-5 Receiving Hearing Screening Tests by Age Group, Tribal Affiliation, Ethnicity, Race and Year


Percent of children receiving hearing screening tests from ages 1-5 by ethnicity




Percent of children receiving hearing screening tests from ages 1-5 by sex



Percent of children receiving hearing screening tests from ages 1-5 by race



Percent of children receiving hearing screening tests from ages 1-5 by tribal affiliation



Percent of children receiving hearing screening tests from ages 1-5 by age group



Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

Note: Data was suppressed for Asian/Pacific Islander and Black.




Figure 32. Percent of AHCCCS Children Ages 1-5 Screened for Hearing and Who Had Additional Audiology Services by Subregion by Year


Regional Range 85.7% 100.0%

Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.


Note: Data was suppressed for Blue Ridge (2019), Concho (2017 and 2019), Holbrook (all years), Snowflake (2017) and Vernon (2017). The high percentages are mostly due to low counts, so interpret carefully.

Figure 33. Percent of AHCCCS Children Ages 1-5 Screened for Hearing and Who Had Additional Audiology Services by Age Group, Tribal Affiliation, Ethnicity, Race and Year


Percent of children screened ages 1-5 who had additional audiology services by tribal affiliation




Percent of children screened ages 1-5 who had additional audiology services by sex



Percent of children screened ages 1-5 who had additional audiology services by ethnicity



Percent of children screened ages 1-5 who had additional audiology services by age group



Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

Note: The high percentages for tribally affiliated children and Hispanic or Latino children are due to low counts, so interpret carefully.

Table 20. Percent of Audiology Service Claims by Provider Type for AHCCCS Children Ages 1-5 Screened for Hearing and Who Had Additional Audiology Services, 2017-2019

	2017		2018		2019	
Provider Type	Claims Count	Percent of Total	Claims Count	Percent of Total	Claims Count	Percent of Total
Audiologist	52	73%	62	78%	48	77%
Physician – MD/DO	19	27%	18	23%	8	13%

ORAL HEALTH

Oral health concerns our teeth, gums, and oral-facial system that includes the ability to smile, speak, chew and other senses. Daily brushing and flossing of our teeth and gums demonstrates good oral hygiene, but it is not enough to maintain good oral health. We also need good nutrition, proper management of other health conditions, access to dental care, and extra help when there is a genetic predisposition to oral health conditions or special health care needs.

Oral health is a key indicator of overall health, well-being and quality of life.

Unfortunately, tooth decay has become a chronic disease in children. The CDC reports that 20% of children ages 5-11 have at least one untreated cavity, and children in low-

income families are twice as likely to have cavities than children in higher-income families (Dye, Xianfen, & Beltrán-Aguilar, 2012). Cavities can be prevented by applying a fluoride varnish to primary and permanent teeth, drinking fluoridated tap water, brushing with a fluoride toothpaste, and applying dental sealants. Children should have regular visits to the dentist, beginning before their first birthday, for early identification and management of problems (Enany, n.d.). This report focuses on dental visits for ages 1-5.

In Navajo/Apache Region, 39-43% of AHCCCS children had at least one annual dental visit compared to 51-53% of AHCCCS children statewide in Table 21. Neither the region nor the state met the AHCCCS MPS of 60% for annual dental visits for ages 2-20 in Table 22. No subregion met the AHCCCS MPS in Figure 34. AHCCCS children ages 3-5 (48-54%) were more likely to have at least one annual dental visit than ages 1-2 (24-27%) in Figure 35; Hispanic or Latino AHCCCS children (41-50%) were more likely to have at least one annual dental visit than Non-Hispanic or Latino AHCCCS children (39-42%).

Two preventative care dental visits are recommended annually for children. Regionally, 12-14% of AHCCCS children received the biannual preventative care dental visit compared to 18-20% of AHCCCS children statewide (Table 21); 36-41% of regional AHCCCS children had at least one preventative care dental visit per year in Figure 36.

In Table 21, fluoride varnish was applied to 34-38% of AHCCCS children in the region compared to 47-49% of AHCCCS children statewide. AHCCCS children ages 3-5 (42-48%) were more likely to have a fluoride

varnish application than ages 1-2 (21-23%) in Figure 38. Hispanic or Latino AHCCCS children (36-46%) were more likely to have a fluoride varnish application than Non-Hispanic or Latino AHCCCS children (34-37%).

Table 21. Percent of Statewide and Regional AHCCCS Children Oral Health Visits for Ages 1-5, 2017-2019

Type of Visit / Year	2017		20	18	2019	
	Region	Arizona	Region	Arizona	Region	Arizona
Any Annual Dental Visit	43%	51%	39%	52%	43%	53%
Preventative Care Dental Visit Twice Annually	14%	18%	12%	19%	13%	20%
Fluoride Varnish Application	37%	47%	34%	48%	38%	49%

Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

Table 22. AHCCCS Statewide Contractor Rate of Performance on Annual Dental Visits for Ages Two to 20 Years, 2017-2019

Contractor	2017	2018	2019	Minimum Performance Standard
AHCCCS Complete Care	61%	61%	60%	60%
Comprehensive Medical and Dental Program	74%	75%	60%	60%
KidsCare	74%	74%	76%	60%

Source: (Health Services Advisory Group, 2021); (Health Services Advisory Group, 2019) (Health Services Advisory Group, 2020).

Table 23. Percent of AHCCCS Claims by Provider Type for Children Ages 1-5 With at Least One Annual Dental Visit, 2017-2019

Provider Type	2017		2018		2019	
	Claims Count	Percent of Total	Claims Count	Percent of Total	Claims Count	Percent of Total
Dentist	2,006	97%	1,817	98%	1,886	99%
Federally Qualified Health Center (FQHC)	24	1%	24	1%	17	1%
Other	34	2%	9	0%	6	<1%




Figure 34. Percent of AHCCCS Children Ages 1-5 With at Least One Annual Dental Visit by Subregion by Year

Regional Range
29.2% 47.1%

Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.


Notes: This indicator includes any claim with an associated dental procedure code (CDT).

Figure 35. Percent of AHCCCS Children Ages 1-5 With at Least One Annual Dental Visit by Age Group, Tribal Affiliation, Ethnicity, Race and Year


Percent of children who have at least one dental visit by age group




Percent of children who have at least one dental visit by tribal affiliation



Percent of children who have at least one dental visit by ethnicity



Percent of children who have at least one dental visit by race



Notes: This indicator includes any claim with an associated dental procedure code (CDT).

100%

80%
60%
40%
39.5%
35.7%
38.8%

2017
2018
2019

Figure 36. Percent of AHCCCS Children Ages 1-5 With One and Two Preventative Care Dental Visits in a Year

Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

Notes: This indicator is called a preventative care dental visit and includes the following procedures: D0120 periodic oral evaluation, D0150 comprehensive oral evaluation and D0145 oral evaluation for patient under 3 years of age and counseling with primary caregiver.

Table 24. Percent of Claims by Provider Type for AHCCCS Children Ages 1-5 Receiving Fluoride Varnish, 2017-2019

	2017		2018		2019	
Provider Type	Claims Count	Percent of Total	Claims Count	Percent of Total	Claims Count	Percent of Total
Dentist	1,243	99%	1,061	97%	1,088	94%
Federally Qualified Health Center (FQHC)	<6	DS	6	1%	8	1%
Physician – MD/DO	9	1%	21	2%	55	5%
Other	<6	DS	7	1%	6	1%




Figure 37. Percent of AHCCCS Children Ages 1-5 Receiving Fluoride Varnish by Subregion by Year


Regional Range 24.6% 41.9%

Figure 38. Percent of AHCCCS Children Ages 1-5 Receiving Fluoride Varnish by Age Group, Tribal Affiliation, Ethnicity, Race and Year


Percent of children receiving flouride varnish ages 1-5 by age group




Percent of children receiving flouride varnish ages 1-5 by ethnicity



Percent of children receiving flouride varnish ages 1-5 by tribal affiliation



Percent of children receiving flouride varnish ages 1-5 by race



IMMUNIZATIONS

Childhood vaccines protect children from many serious and potentially life-threatening diseases such as diphtheria, measles, meningitis, polio, tetanus and whooping cough, at a time in their lives when they are most vulnerable to disease. Approximately 300 children in the United States die each year from vaccine preventable diseases (HHS Office of Disease Prevention and Health Promotion, 2021). Immunizations are essential for disease prevention and are a critical aspect of preventable care for children. Vaccination coverage must be maintained to prevent a resurgence of vaccine-preventable diseases.

The Centers for Medicare and Medicaid Services measures the quality of immunizations through a core indicator of childhood immunization status which is also used by HEDIS. The measure calculates a rate for certain vaccines recommended by a child's second birthday (National Quality Forum, 2017):

- Percent of children who have completed the following schedules: four diphtheria, tetanus and acellular pertussis (DTaP); three polio (IPV); one measles, mumps and rubella (MMR); three haemophilus influenza type B (HiB); three hepatitis B (HepB); one chicken pox (VZV); four pneumococcal conjugate (PCV); one hepatitis A (HepA); two or three rotavirus (RV); two influenza (flu).
- Percent of children who have completed all vaccine courses combined: Combo 10.
- Percent of children who have completed Combo 3: four diphtheria, tetanus and acellular
 pertussis (DTaP); three polio (IPV); one measles, mumps and rubella (MMR); three haemophilus
 influenza type B (HiB); three hepatitis B (HepB); one chicken pox (VZV); four pneumococcal
 conjugate (PCV).

AHCCCS measures childhood immunization completion rates with each of its contractors biennially using the core measure. AHCCCS children's immunization status in Table 26 is the percent of AHCCCS children who have completed each indicated vaccine course by their second birthday, recorded in AHCCCS claims only. These rates were substantially lower than AHCCCS' published statistics in Table 25 due to the limitation of claims data and should be interpreted with caution. AHCCCS declared that claims data does not have the greatest level of detail as claims are not always reported for immunizations, particularly in school settings. To accurately capture immunization rates in AHCCCS' published statistics, AHCCCS uses data from medical records and from the Arizona State Immunization Information System (ASIIS), which is maintained by the Arizona Department of Health Services.

AHCCCS reported that statewide childhood immunization completion rates met or exceeded the national mean rates for three immunizations: DTaP, Hep A and Combo 3 (Arizona Health Care Cost Containment System, 2018) (Table 25). Several barriers to immunizations remained, such as the spread of misinformation about vaccines and parental hesitancy. The rate of exemptions from immunizations increased statewide as nearly 6% of kindergarteners had a Personal Beliefs Exemption in place since the 2017-2018 school year (Arizona Department of Health Services, 2021).

Table 25. AHCCCS Statewide Aggregate Immunization Completion Rates by Two Years Old, FFY 2016

Immunizations	FFY 2016 (period ending 9/30/2017)	HEDIS Medicaid Mean	AHCCCS Minimum Performance Standard
DTaP	79%	77%	85%
Polio	88%	89%	91%
MMR	89%	90%	91%
HiB	87%	88%	90%
Нер В	87%	88%	90%
VZV	88%	89%	88%
PCV	76%	77%	82%
Нер А	88%	84%	40%
RV	61%	69%	60%
Flu	40%	45%	45%
Combo 3	71%	70%	68%

Source: (Arizona Health Care Cost Containment System, 2018).

Note: The rows shaded green are the childhood immunization rates that met or exceeded the national median rates.

Table 26. Percent of Statewide and Regional AHCCCS Children Immunization Status, from AHCCCS Claims Data Only, 2017-2019

	20	17	20	018	20)19
Immunizations	Region	Arizona	Region	Arizona	Region	Arizona
DTaP	15%	30%	25%	38%	39%	52%
Polio	20%	40%	33%	51%	51%	66%
MMR	41%	57%	62%	72%	64%	76%
HiB	21%	44%	39%	56%	47%	69%
Нер В	7%	13%	12%	18%	19%	21%
VZV	41%	57%	60%	72%	62%	76%
PCV	3%	18%	18%	31%	35%	52%
Нер А	45%	65%	55%	75%	57%	78%
RV	11%	31%	18%	39%	35%	51%
Flu	8%	19%	16%	31%	19%	34%
Combo 3	<1%	4%	5%	10%	9%	15%
Combo 10	DS	2%	1%	4%	4%	7%

Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

Notes: Rates only include immunizations recorded in AHCCCS claims, this is likely an undercount of immunization rates. Rows shaded pink are for comparing with Table 25.

MATERNAL PRENATAL AND POSTPARTUM CARE

Research has shown that the health of women before pregnancy and after delivery significantly impacts the health of their babies; therefore, it is important to focus on women's preconception health, prenatal care, postpartum care and beyond (Healthy People 2030).

Women who do not seek prenatal care are three times as likely to deliver a low birth weight infant.

(NICHD - Eunice Kennedy Shriver National Institute of Child Health and Human Development, 2017) For pregnant women, prenatal care is essential for a healthy pregnancy and reducing the complications that can lead to poor birth outcomes for mother and child. Prenatal care involves regular visits to a health care provider to monitor the mother's health and health of the developing fetus, and this care should begin as

early as possible in the pregnancy and continue until delivery. Prenatal care can identify problems or complications and take steps to manage them (NICHD - Eunice Kennedy Shriver National Institute of Child Health and Human Development, 2017). The American Academy of Pediatrics and the American College of Obstetricians and Gynecologists recommend that a woman with an uncomplicated pregnancy be examined at least once in the first trimester for prenatal care. Appropriate perinatal services and education are crucial components of a healthy birth.

The period of up to 60 days following childbirth is called the postpartum period. Preexisting health conditions, social determinants, and newly developed conditions contribute to maternal morbidity and mortality during this period. Health care providers consider the postpartum period to be critical to the health and well-being of both mother and child, so postpartum care should not be considered as optional. Yet, research has shown that nearly 40% of women in the United States have gone without a single postpartum visit (American College of Obstetricians and Gynecologists, 2018).

In Navajo/Apache Region, 92-95% of pregnant women began prenatal care in the first trimester compared to 84-86% of AHCCCS women statewide in Table 27, which are both above the Healthy People 2030 target rate of 81%²². All subregions met or exceeded the Healthy People 2030 target rate for timely prenatal care for all years, except for Holbrook which exceeded the target rate in 2019 only (Figure 39). There was little difference between AHCCCS women's age, tribal affiliation, race or ethnicity in receiving timely prenatal care as all had high rates for this indicator in Figure 40.

For postpartum care, 96% of regional AHCCCS women had at least one postpartum visit compared to 88-89% of AHCCCS women statewide (Table 27) and 64-75% of Medicaid women nationally. In Figure 41, all subregions exceeded the state rates for postpartum care in all years, except for Holbrook which exceeded the state rate in 2019 only.

²² Healthy People 2030 Prenatal Care Objective - https://health.gov/healthypeople/objectives-and-data/browse-objectives/pregnancy-and-childbirth/increase-proportion-pregnant-women-who-receive-early-and-adequate-prenatal-care-mich-08


Table 27. Percent of All AHCCCS Women Who Received Timely Prenatal and Postpartum Care, 2017-2019

Type of Care	2017		20	18	2019	
Type of Care	Region	Arizona	Region	Arizona	Region	Arizona
Prenatal Care	92%	84%	95%	86%	93%	85%
Postpartum Care	96%	88%	96%	89%	96%	89%

Figure 39. Percent of Deliveries That Received a Prenatal Care Visit While Enrolled in AHCCCS in the First Trimester, on the Enrollment Start

Date or Within 42 Days of Enrollment in AHCCCS by Subregion by Year


2017 2018 2019




Regional Range 78.0% 100.0%

Figure 40. Percent of Deliveries That Received a Prenatal Care Visit While Enrolled in AHCCCS in the First Trimester, on the Enrollment Start Date or Within 42 Days of Enrollment in AHCCCS by Age Group, Tribal Affiliation, Ethnicity, Race and Year


Percent of deliveries that received a prenatal care by age group




Percent of deliveries that received a prenatal care by ethnicity



Percent of deliveries that received a prenatal care by tribal affiliation



Percent of deliveries that received a prenatal care by race



Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

Note: Data was suppressed for Asian/Pacific Islander (all years and Black (2018). The high percentages for Black are due to low counts.





Figure 41. Percent of AHCCCS Deliveries That Had a Postpartum Visit After Delivery by Subregion by Year


Regional Range 82.9% 100.0%

Figure 42. Percent of AHCCCS Deliveries That Had a Postpartum Visit After Delivery by Age Group, Race, Ethnicity, Tribal Affiliation and Year


Percent of deliveries that had a postpartum visit after delivery by age group




Percent of deliveries that had a postpartum visit by tribal affiliation



Percent of deliveries that had a postpartum visit by ethnicity



Percent of deliveries that had a postpartum visit by race



Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

Note: Data was suppressed for Asian/Pacific Islander (all years and Black (2018). The high percentages for Black are due to low counts.

HEALTH PLAN PERFORMANCE

This section provided a selection of health indicators to compare results among the AHCCCS health plans.

Table 29 provided a snapshot of service utilization for regional AHCCCS children enrolled in each AHCCCS health plan. The following regional health plans met or exceeded the AHCCCS statewide aggregate performance and MPS for PCP visits for ages 1-5: AZ Complete Health (2017 and 2018), Banner University Family Care (2019), Comprehensive Medical and Dental Program (CMDP) (2018 and 2019), Mercy Care Plan (2018 and 2019) and UnitedHealthcare (2017 and 2019). For at least one well-child visit in the first 15 months, all regional health plans performed well (86-100%), except for the American Indian Health Program. For six or more well-child visits in the first 15 month, only CMDP exceeded the AHCCCS statewide aggregate performance and MPS in 2018. For one or more well-child visits for ages 3-5, four health plans met or exceeded the AHCCCS statewide aggregate performance and MPS: Banner University Family Care (2019), Children's Rehabilitative Services (2018), Mercy Care Plan (2018) and UnitedHealthcare (2019). Only Banner University Family Care (2019) met or exceeded the AHCCCS statewide aggregate health plan performance or MPS for preventative care dental visits for ages 1-5.²³

_

²³ The AHCCCS statewide indicator for preventative care dental visits includes ages 2-20 which incorporates a significantly larger number of AHCCCS children than our reporting on ages birth to 5, so the rates should be compared with caution.

Table 28. AHCCCS Statewide Aggregate Health Plan Performance and Minimum Performance Standards, 2017-2019

Year/Indicator	Health Plan Type	One or More Annual PCP Visits, Ages 1-6 (MPS)	Six or More Well-Child Visits in First 15 Months (MPS)	One or More Annual Well- Child Visits, Ages 3-6 (MPS)	One or More Annual Preventative Care Dental Visits, Ages 2-20 (MPS)
	Acute	83% (84%)	60% (65%)	61% (66%)	61% (60%)
2017	CMDP	92% (84%)	75% (65%)	75% (66%)	74% (60%)
	CRS	93% (84%)	49% (65%)	66% (66%)	67% (60%)
	Acute	84% (84%)	62% (65%)	61% (66%)	61% (60%)
2018	CMDP	93% (84%)	N/A (65%)	73% (66%)	75% (60%)
	CRS	92% (84%)	47% (65%)	64% (66%)	68% (60%)
	Acute	84% (87%)	64% (62%)	63% (66%)	60% (60%)
2019	CMDP	92% (87%)	N/A (62%)	75% (72%)	75% (60%)
	CRS	N/A (87%)	N/A (62%)	N/A (66%)	N/A (60%)

Source: (Health Services Advisory Group, 2019); (Health Services Advisory Group, 2020); (Health Services Advisory Group, 2021); (Health Services Advisory Group, 2019); (Health Services Advisory Group, 2020).

Notes: There was no MPS for DES/DDD for six or more well-child visits in the first 15 months. Cells for which data was not available are indicated by "N/A".

Table 29. Select Regional Indicators by AHCCCS Health Plan, 2017-2019

		One or More	Well-Child Visits in First 15 Months		One or More	One or More Preventative	
Health Plan	Year	PCP Visits, Ages 1-5	At Least One Visit*	Six or More Visits	Well-Child Visits, Ages 3-5	Care Dental Visits, Ages 1-5	
ALICCOS Amenican Indian	2017	13%	36%	DS	6%	14%	
AHCCCS American Indian Health Program	2018	7%	DS	DS	DS	17%	
Ticalti i Togram	2019	19%	DS	DS	11%	22%	
	2017	86%	86%	DS	DS	29%	
AZ Complete Health	2018	92%	DS	DS	DS	46%	
	2019	85%	DS	DS	52%	58%	
B	2017	68%	100%	DS	47%	36%	
Banner University Family Care	2018	83%	DS	DS	50%	26%	
Care	2019	94%	DS	DS	82%	65%	
	2017	82%	100%	DS	DS	38%	
Care 1st	2018	74%	DS	DS	DS	DS	
	2019	82%	95%	52%	53%	33%	
Children's Rehabilitative	2017	89%	100%	DS	61%	53%	
Services (CRS)	2018	91%	DS	DS	69%	56%	
DES Comprehensive	2017	81%	92%	DS	61%	36%	
Medical and Dental	2018	89%	90%	70%	66%	40%	
(CMDP) Program	2019	90%	100%	DS	55%	54%	
	2017	81%	100%	60%	27%	35%	
Mercy Care Plan	2018	93%	89%	DS	63%	42%	
	2019	87%	100%	55%	40%	48%	
Canada Harlah Chaire	2017	81%	90%	47%	53%	47%	
Steward Health Choice	2018	81%	90%	46%	53%	40%	
AZ	2019	83%	91%	54%	49%	43%	
	2017	84%	93%	47%	53%	48%	
United Healthcare	2018	81%	93%	39%	55%	48%	
	2019	87%	100%	DS	74%	50%	

Source: AHCCCS Claims Data, 2021. CHiR is the source for all processing of the AHCCCS data.

Notes: Cells with "DS" represented suppressed data due to low counts. Cells shaded green indicate the rate met or exceeded the AHCCCS statewide performance by health plan (Table 30); cells shaded blue indicate the health plan met or exceeded the AHCCCS statewide aggregate performance, cells shaded pink indicate the health plan met or exceeded the MPS, and cells shaded purple indicate the health plan met or exceeded both the MPS and statewide aggregate performance (see Table 28).

Table 30. Select AHCCCS Statewide Indicators by Health Plan, 2017-2019

Health Plan	Year	One or More PCP Visits, Ages 1-6	Well-Child Visits in First 15 Months Six or More Visits	One or More Well- Child Visits, Ages 3-6	One or More Preventative Care Dental Visits, Ages 2-20
	2017	N/A	N/A	N/A	N/A
AZ Complete Health	2018	82%	61%	59%	48%
	2019	82%	64%	61%	55%
	2017	N/A	N/A	N/A	N/A
Banner University Family Care	2018	84%	62%	60%	54%
	2019	83%	64%	61%	53%
	2017	83%	66%	64%	62%
Care 1st	2018	86%	67%	67%	65%
	2019	84%	71%	64%	63%
	2017	N/A	N/A	N/A	N/A
Magellan Complete Care	2018	N/A	N/A	N/A	N/A
	2019	67%	N/A	47%	37%
	2017	85%	63%	62%	64%
Mercy Care Plan	2018	86%	66%	63%	64%
	2019	87%	65%	65%	63%
	2017	83%	59%	60%	61%
United Healthcare	2018	84%	61%	61%	62%
	2019	86%	66%	67%	62%

Source: (Health Services Advisory Group, 2019); (Health Services Advisory Group, 2020); (Health Services Advisory Group, 2021); (Health Services Advisory Group, 2019); (Health Services Advisory Group, 2020).

Notes: Cells that did not have available data for that year and/or the health plan was not contracted for that year are indicated by "N/A". Cells shaded blue indicate the rate met or exceeded the AHCCCS statewide aggregate health plan performance rate; cells shaded purple indicate the rate met or exceeded both the AHCCCS statewide aggregate health plan performance rate and the AHCCCS MPS (see Table 28). AHCCCS did not measure health plan performance for the indicator of at least one PCP visit by 15 months of age.

CONCLUSION

The physical, mental, and emotional health of young children lays the foundation for the rest of their life. Navajo/Apache Region had several assets contributing to better health outcomes for young children and women enrolled in AHCCCS from 2017 to 2019, including newborn hearing screenings and prenatal and postpartum care. These achievements contributed to good health outcomes throughout the region. The areas where needs were identified for AHCCCS women and children included PCP and well-child visits, the supply of health professionals, developmental screening, lead poisoning screening, vision screening and oral health.

The information in this report can be combined with other available information to create a more comprehensive view of young children and women in the region for regional council planning.

REFERENCES

- American Academy of Child and Adolescent Psychiatry. (2006). The Early Childhood Service Intensity System (ECSII). Retrieved from Arizona Health Care Cost Containment System:

 https://www.azahcccs.gov/PlansProviders/Downloads/TI/CoreComponents/The%20Early%20Childhood%20Service%20Intensity%20Instrument%20(ESCII)%20.pdf
- American Board of Pediatrics. (2018). Pediatric Physicians Workforce Data Book, 2017-2018. Chapel Hill, NC: American Board of Pediatrics. Retrieved from https://www.abp.org/sites/abp/files/pdf/pediatricphysiciansworkforcedatabook2017-2018.pdf
- American Board of Pediatrics. (2019). Pediatric Physicians Workforce Data Book, 2018-2019. Chapel Hill, NC: American Board of Pediatrics. Retrieved from https://www.abp.org/sites/abp/files/workforcedata2018-2019.pdf
- American Board of Pediatrics. (2020). Pediatric Physicians Workforce Data Book, 2019-2020. Chapel Hill, NC: American Board of Pediatrics. Retrieved from https://www.abp.org/sites/abp/files/pdf/workforcedata2019-2020.pdf
- American College of Obstetricians and Gynecologists. (2018). Optimizing postpartum care. ACOG Committee Opinion No. 736. Obstet Gynecol, 131(e140-150). Retrieved from https://www.acog.org/-/media/project/acog/acogorg/clinical/files/committee-opinion/articles/2018/05/optimizing-postpartum-care.pdf
- American Dental Association. (2021). Supply of Dentists in the U.S.: 2001 2020. Chicago: Health Policy Institute, American Dental Association. Retrieved from https://www.ada.org/~/media/ADA/Science%20and%20Research/HPI/Files/HPIData_SOD_2020.xlsx?la=en
- American Public Health Association. (2019). Coordinated nationwide approaches to promote eye health and reduce vision impairment. Policy No 20191. Retrieved September 8, 2021, from https://www.apha.org/policies-and-advocacy/public-health-policy-statements/policy-database/2020/01/10/coordinated-nationwide-approaches-to-promote-eye-health-and-reduce-vision-impairment
- Arizona Department of Health Services. (2003, July 8). Lead A Silent Poison. Retrieved from Arizona Department of Health Services:

 https://www.azdhs.gov/documents/preparedness/epidemiology-disease-control/lead-poisoning/lead-a-silent-poison-brochure.pdf
- Arizona Department of Health Services. (2006, August 1). What everyone should know about Lead Poisoning. Retrieved from Arizona Department of Health Services:

 https://www.azdhs.gov/documents/preparedness/epidemiology-disease-control/lead-poisoning/lead-poisoning-brochure-english.pdf
- Arizona Department of Health Services. (2021). Arizona 2019 Annual Report Blood Lead Surveillance Childhood Lead Poisoning Prevention Program. Phoenix: Office of Environmental Health, Bureau

- of Epidemiology and Disease Control . Retrieved October 11, 2021, from https://azdhs.gov/documents/preparedness/epidemiology-disease-control/lead-poisoning/2019-annual-blood-lead-report.pdf
- Arizona Department of Health Services. (2021). Arizona Immunization Program. Retrieved August 30, 2021, from https://www.azdhs.gov/preparedness/epidemiology-disease-control/immunization/index.php#reports-immunization-coverage
- Arizona Department of Health Services. (2021, September). Consumers Provider & Facility Databases.

 Retrieved from Medical Facilities Licensing: https://azdhs.gov/licensing/medical-facilities/index.php#consumers-databases
- Arizona Department of Health Services. (n.d.). Newborn Hearing Screening Early Hearing Detection and Intervention 101. Retrieved from Arizona Department of Health Services: https://azdhs.gov/preparedness/state-laboratory/newborn-screening/index.php#az-ehdi-home
- Arizona Health Care Cost Containment System. (2018). AHCCCS Complete Care: The Future of Integrated Healthcare Delivery. Retrieved from AHCCCS: https://azahcccs.gov/AHCCCS/Initiatives/AHCCCSCompleteCare/
- Arizona Health Care Cost Containment System. (2018). Childhood Immunization Completion Rates.

 Division of Health Care Management, Phoenix. Retrieved from

 https://www.azahcccs.gov/Resources/Downloads/PerformanceMeasures/CYE2017_ChildhoodI

 mmunizationReport.pdf
- Arizona Health Care Cost Containment System. (2018, August 8). Newborn Screening. Phoenix, AZ.

 Retrieved from

 https://www.azahcccs.gov/Resources/Downloads/Training/Back_To_Basics_Newborn_Screening.pdf
- Arizona Health Care Cost Containment System. (2018, January 18). Working with birth through five population practice tool. Retrieved August 30, 2021, from https://www.azahcccs.gov/PlansProviders/Downloads/GM/ClinicalGuidanceTools/BirthThrough Five/WorkingWithTheBirthThroughFivePopulationPracticeTool.pdf
- Arizona Health Care Cost Containment System. (2021). Performance improvement project developmental screening interim report. Phoenix. Retrieved from https://www.azahcccs.gov/resources/Downloads/PerformanceImprovementProjects/AHCCCSD evelopmentalScreeningPIPInterimReport.pdf
- Arizona Health Care Cost Containment System. (2021, October). Policy 430 Attachment A AHCCCS Early and Periodic Screening, Diagnostic and Treatment Periodicity Schedule. Retrieved from AHCCCS Medical Policy Manual: https://www.google.com/url?client=internal-element-cse&cx=016416319741515762556:nfpz2wqmsus&q=https://www.azahcccs.gov/shared/Downlo ads/MedicalPolicyManual/400/430_AttachmentA.docx&sa=U&ved=2ahUKEwjs5f_b8rL3AhWsJUQIHf4wCCIQFnoECAMQAQ&usg=AOvVaw2PugsSuiAzf

- Arizona Health Care Cost Containment System. (July 2021). AHCCCS Quality Strategy Evaluation CYE 2018 CYE 2020. Phoenix. Retrieved from https://azahcccs.gov/PlansProviders/Downloads/QualityStrategyEvaluationCYE2018-CYE2020.pdf
- Association of American Medical Colleges. (2017). 2017 State Physician Workforce Data Report. Washington, DC: AAMC.
- Association of American Medical Colleges. (2019). 2019 State Physician Workforce Data Report. Washington, DC: AAMC.
- Centers for Disease Control and Prevention. (n.d.). Adverse Childhood Experiences. (N. C. Control, Producer) Retrieved from Violence Prevention:

 https://www.cdc.gov/violenceprevention/aces/index.html
- Centers for Disease Control and Prevention. (n.d.). Learn more about your child's development: developmental monitoring and screening. Retrieved August 27, 2021, from https://www.cdc.gov/ncbddd/actearly/pdf/Dev-Mon-and-Screen-English-and-Spanish-P.pdf
- Child and Adolescent Health Measurement Initiative. (n.d.). 2019-2020 National Survey of Children's Health (NSCH) data query. Data Resource Center for Child and Adolescent Health supported by the U.S. Department of Health and Human Services, Health Resources and Services Administration (HRSA), Maternal and Child Health Bureau (MCHB). Retrieved May 3, 2022, from https://www.childhealthdata.org/
- Choo, Y. Y., Agarwal, P., How, C. H., & Yeleswarapu, S. P. (2019). Developmental delay: identification and management at primary care level. Singapore Medical Journal, 60(3), 119-123. doi:10.11622/smedj.2019025
- Dye, B. A., Xianfen, L., & Beltrán-Aguilar, E. D. (2012). Selected oral health indicators in the United States 2005–2008. Hyattsville, MD: National Center for Health Statistics, Centers for Disease Control and Prevention.
- Enany, E. (n.d.). Small smiles: what you need to know about baby teeth. (A. D. Association, Producer)

 Retrieved August 28, 2021, from mouthhealthy.org: https://www.mouthhealthy.org/en/babies-and-kids/childrens-dental-health
- Eyes on Learning. (n.d.). Why Screening. Retrieved from Eyes on Learning: http://eyesonlearning.org/why-childrens-vision-matters/why-screening/
- Hales, C. M., Carroll, M. D., Fryar, C. D., & Ogden, C. L. (2017). Prevalence of obesity among adults and youth: United States, 2015–2016. Hyattsville, MD:: National Center for Health Statistics.
- Health Services Advisory Group. (2019). CYE 2018 Annual Report for Acute Care and CMDP. Phoenix:

 Arizona Health Care Cost Containment System. Retrieved from

 https://www.azahcccs.gov/Resources/Downloads/EQR/2018/CYE_2018_External_Quality_Revie

 w_Annual_Report_Acute_Care_And_CMDP.pdf

nual Report ALTCS.pdf

- Health Services Advisory Group. (2019). External Quality Review Annual Report for Arizona Long Term
 Care System Contractors. Phoenix: Arizona Health Care Cost Containment System. Retrieved
 from
 https://azahcccs.gov/Resources/Downloads/EQR/2018/CYE_2018_External_Quality_Review_An
- Health Services Advisory Group. (2020). CYE 2019 Annual Report for 2018 Acute Care and CMDP, RBHAs, and CRS. Phoenix: Arizona Health Care Cost Containment System. Retrieved from https://www.azahcccs.gov/Resources/Downloads/EQR/2019/CYE2019ExternalQualityReviewAn nualReport-AcuteCMDPRBHACRS.pdf
- Health Services Advisory Group. (2020). External Quality Review Annual Report for Arizona Long Term Care System Contractors. Phoenix: Arizona Health Care Cost Containment System. Retrieved from https://azahcccs.gov/Resources/Downloads/EQR/2019/CYE2019ExternalQualityReviewAnnualReportALTCS.pdf
- Health Services Advisory Group. (2021). CYE 2019 Annual Report for AHCCCS Complete Care and CMDP. Phoenix: Arizona Health Care Cost Containment System. Retrieved October 18, 2021, from https://www.azahcccs.gov/Resources/Downloads/EQR/2020/CYE2020ExternalQualityReviewAn nualReportACCandCMDP.pdf
- Health Services Advisory Group. (2021). External Quality Review Annual Report for Arizona Long Term Care System. Phoenix: Arizona Health Care Cost Containment System. Retrieved from https://azahcccs.gov/Resources/Downloads/EQR/2020/CYE2020ExternalQualityReviewAnnualReportEPDandDDD.pdf
- Healthy People 2030. (n.d.). Pregnancy and Childbirth. Office of Disease Prevention and Health Promotion, Office of the Assistant Secretary for Health, Office of the Secretary, U.S. Department of Health and Human Services. Retrieved August 25, 2021, from https://health.gov/healthypeople/objectives-and-data/browse-objectives/pregnancy-and-childbirth
- HHS Office of Disease Prevention and Health Promotion. (2021, October). Immunization and Infectious Diseases. Retrieved from HealthyPeople.gov: https://www.healthypeople.gov/node/3527/data-details.%C2%A0Accessed
- Moreno, M. A. (2018). The Well-Child Visit. JAMA Pediatr, 172(1), 104. doi:10.1001/jamapediatrics.2017.4041
- National Center for Environmental Health. (2020, January). Health Effects of Lead Exposure. Retrieved from Centers for Disease Control and Prevention:

 https://www.cdc.gov/nceh/lead/prevention/health-effects.htm
- National Center on Birth Defects and Developmental Disabilities. (2019, August). Summary of 2017

 National CDC EHDI Data. Retrieved from Centers for Disease Control and Prevention:

- https://www.cdc.gov/ncbddd/hearingloss/2017-data/documents/01-2017-HSFS-Data-Summary.pdf
- National Committee for Quality Assurance. (2021). Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents. Retrieved from HEDIS:

 https://www.ncqa.org/hedis/measures/weight-assessment-and-counseling-for-nutrition-and-physical-activity-for-children-adolescents/
- National Council of State Boards of Nursing. (2021). Number of Nurses in U.S. and by Jurisdiction. Retrieved from NCSBN: https://www.ncsbn.org/14283.htm
- National Quality Forum. (2017, January 23). Childhood Immunization Status. Retrieved August 30, 2021, from https://www.qualityforum.org/QPS/QPSTool.aspx?m=240#qpsPageState=%7B%22TabType%22%3A1,%22TabContentType%22%3A2,%22ItemsToCompare%22%3A%5B%5D,%22StandardID%22%3A240,%22EntityTypeID%22%3A1%7D
- National Scientific Council on the Developing Child. (2004). Children's Emotional Development Is Built into the Architecture of Their Brains. Center on the Developing Child at Harvard University.

 Retrieved from http://www.developingchild.net
- NICHD Eunice Kennedy Shriver National Institute of Child Health and Human Development. (2017, January 31). Pre-Pregnancy Care and Prenatal Care. Retrieved August 27, 2021, from https://www.nichd.nih.gov/health/topics/factsheets/preconceptioncare
- NIDCD Information Clearinghouse. (2011, May). NIDCD Fact Sheet: It's Important to Have Your Baby's Hearing Screened. Retrieved from National Institute on Deafness and Other Communication: https://www.nidcd.nih.gov/sites/default/files/Documents/health/hearing/ItsImportantToHaveY ourBabysHearingScreenedWebFS.pdf
- Piehl, M. D., Clemens, C. J., & Joines, J. D. (2000, Aug). "Narrowing the Gap": decreasing emergency department use by children enrolled in the Medicaid program by improving access to primary care. Arch Pediatr Adolesc Med, 154(8), 791-5. doi:10.1001/archpedi.154.8.791
- Sturgeon, M. (2015, December 15). Parent Plus: Importance of routine pediatrician visits. AAP News. The American Academy of Pediatrics. Retrieved from https://www.aappublications.org/news/aapnewsmag/2015/12/15/WellChild121515.full.pdf
- Transforming Clinical Practice Initiative. (2019, December 6). Reducing Unnecessary Emergency Department Visits. Retrieved from Centers for Medicare and Medicaid Services: https://innovation.cms.gov/files/x/tcpi-changepkgmod-edvisits.pdf
- Zablotsky, B., & Black, L. (2020). Prevalence of children aged 3–17 years with developmental disabilities, by urbanicity: United States, 2015–2018. Hyattsville, MD: National Health Statistics Reports No 139. Retrieved from https://www.cdc.gov/nchs/data/nhsr/nhsr139-508.pdf

APPENDIX: DATA SOURCES

The source of data in all tables, graphs, reports, presentations, and other publications is the Arizona Health Care Cost Containment System (AHCCCS) (2021) unless otherwise noted. CHiR is the source of the calculations, analysis and/or processing of the data.

AHCCCS health claims and encounters data. AHCCCS is the state Medicaid provider. The data include health care transactions (paid claims) on all members, patients receiving inpatient, emergency department or other outpatient care in the state. The data layout is already at the individual patient level when received and requires no further manipulation to standardize variables or match patients.

AHCCCS System

In October 2018, AHCCCS enacted major changes to its care delivery system to integrate physical and behavioral health care under designated health plans for its eligible populations, called AHCCCS Complete Care. Integrated care would result in better coordination among providers in the same network and better health outcomes for AHCCCS enrollees. Under AHCCCS Complete Care, the choice of health plans varies by geographic area, but affected members have the same array of covered services and access to a network of providers (Arizona Health Care Cost Containment System, 2018).

Table 31. AHCCCS Complete Care Health Plans by Geographic Service Area

Geographic Service Area	AHCCCS Complete Care Health Plans
North (Apache, Coconino, Mohave, Navajo and Yavapai Counties)	Care 1st and Health Choice Arizona
Central (Maricopa, Gila and Pinal Counties)	Banner University Family Care, Care 1st, Health Choice Arizona, Arizona Complete Health, Magellan Complete Care, Mercy Care, UnitedHealthcare Community Plan
South (Cochise, Graham, Greenlee, La Paz, Pima, Santa Cruz and Yuma Counties)*	Banner University Family Care, Arizona Complete Health, UnitedHealthcare Community Plan (Pima County only)

^{*}Zip codes 85542, 85192, 85550 are in the South geographic service area.

Other health plans serve specialty populations. AHCCCS members with developmental disabilities who are enrolled in the Department of Economic Security/Division of Developmental Disabilities (DES/DDD) with a Children's Rehabilitative Services designation receive integrated care through their assigned DDD health plan. Arizona Long Term Care members receive services through program contractors.

American Indian members have the choice of enrolling in an AHCCCS Complete Care managed care plan or the American Indian Health Program (AIHP fee-for-service) for integrated care or switch enrollment between the two at any time. AIHP members can also choose care coordination through a Tribal Regional Behavioral Health Authority when available (secondary health plan). American Indian members can receive services at any time from an Indian Health Service facility, or a tribally owned or operated organization (i.e., Tribal 638 providers or Urban Indian Health providers).

American Indian members determined to have a Serious Mental Illness receive behavioral health services from a Regional Behavioral Health Authority but have the option to choose the American Indian Health Program or AHCCCS Complete Care for physical health services.

Arizona Health Care Workforce- Physicians. For the provider indicators, we capture the supply of Arizona physicians by specialty using the Arizona Health Care Workforce data set. This data set includes administrative data collected from the Arizona Medical Board and the Arizona Board of Osteopathic Examiners in Medicine and Surgery, the licensing agencies for physicians.